

PROPOSAL FOR EXPERIMENT AT RCNP

28 Jan 2000

TITLE: Neutron and γ Decay from Hole States in Light Nuclei**SPOKESPERSONS:**

Masaru Yosoi

Institution Research Center for Nuclear Physics, Osaka University

Phone/FAX number +81-6-6789-8932 / +81-6-6789-8899

E-mail yosoi@rcnp.osaka-u.ac.jp

Hidenori Toyokawa

Institution Japan Synchrotron Radiation Research Institute (JASRI)

Phone/FAX number +81-791-58-1842 / +81-791-58-0830

E-mail toyokawa@spring8.or.jp

EXPERIMENTAL GROUP:

Name	Institution	Title or Position
Mamoru Fujiwara	RCNP, Osaka University	(AP)
Testuo Noro	RCNP, Osaka University	(AP)
Hisako Fujimura	RCNP, Osaka University	(D3)
Hidetomo Yoshida	RCNP, Osaka University	(D2)
Emi Obayashi	RCNP, Osaka University	(D1)
Harutaka Sakaguchi	Department of Physics, Kyoto University	(AP)
Masanobu Nakamura	Department of Physics, Kyoto University	(L)
Hiroyuki Takeda	Department of Physics, Kyoto University	(D3)
Masatoshi Itoh	Department of Physics, Kyoto University	(D3)
Takahiro Kawabata	Department of Physics, Kyoto University	(D2)
Takatsugu Ishikawa	Department of Physics, Kyoto University	(D1)
Makoto Uchida	Department of Physics, Kyoto University	(M2)
Yuusuke Yasuda	Department of Physics, Kyoto University	(M1)
Hidetoshi Akimune	Department of Physics, Konan University	(L)
Kaoru Yamasaki	Department of Physics, Konan University	(M2)
Yoshitaka Itow	ICRR, University of Tokyo	(A)
Kazuyoshi Kobayashi	ICRR, University of Tokyo	(D1)
Hiroyasu Ejiri	Nuclear Physics Lab., University of Washington	(P)
Taiichi Yamada	Kanto Gakuin University	(AP)

RUNNING TIME:

Installation time without beam	3 days
Test running time for experiment	2 days
Data runs	8 days

BEAM LINE:

Ring : WS course

BEAM REQUIREMENTS:

Type of particle	unpolarized p
Beam energy	392 MeV
Beam intensity	≥ 20 nA
Any other requirements	energy resolution ≤ 300 keV halo-free, small emittance

BUDGET:

Experimental expenses	2,800,000 yen
Travel plans	14 participants should be supported by RCNP

TITLE: Neutron and γ Decay from Hole States in Light Nuclei

SPOKESPERSONS: Masaru Yosoi and Hidenori Toyokawa

SUMMARY OF THE PROPOSAL

In order to understand microscopic structures of deep-hole states and their fragmentation mechanism, we have measured the $^{12}\text{C}(p, 2p)^{11}\text{B}^*$ and $^{16}\text{O}(p, 2p)^{15}\text{N}^*$ reaction at $E_p = 392$ MeV together with decay charged particles from s -hole states in ^{11}B and ^{15}N under the previous E81 and E110 experiments. Our results can not be reproduced by the statistical model calculation using the code CASCADE but are qualitatively consistent with the calculation in the microscopic cluster model with the $SU(3)(\lambda\mu) = (04)$ wave function for the $^{11}\text{B}(s\text{-hole})$ and the $SU(3)(\lambda\mu) = (00)$ for $^{15}\text{N}(s\text{-hole})$.

A recent shell model calculation shows that the s -hole states split into two or more components. In the case of $^{11}\text{B}(s\text{-hole})$, some bump structures are observed in our experimental excitation spectrum, while those are not clear for the $^{15}\text{N}(s\text{-hole})$. As each bump has approximately a different configuration, e.g., $SU(3)[4421](04)$ or $SU(3)[443](04)$ component for the $^{11}\text{B}(s\text{-hole})$, it is predicted that the relative ratio of decay particles is different in each bump. The branching ratio of neutron decay in different excitation energy region is one of the key points to study those bump structures as well as it makes the measurement of fragmentation of deep-hole states complete. Whether the particles decay to the ground states or to excited states of the residual nuclei is another interesting issue. In our E110 results, the decay to ground state is dominant for each particle in the $^{11}\text{B}(s\text{-hole})$ case, while the decay to excited states are mainly occurred in the $^{15}\text{N}(s\text{-hole})$. However, experimental decay ratios between the ground state and excited states have suffered large effects due to the rather high threshold of the detection energy. Measurements of deexcitations from the residual daughter nuclei help to reduce above ambiguities.

In the present proposal, we would like to measure neutron and γ decays from the proton s -hole state in ^{11}B and ^{15}N . The γ decay of ^{15}N s -hole state is also related with the nucleon decay measurement using water Čerenkov detectors. The probability of the high-energy (≥ 15 MeV) γ decay can especially give useful information for the search of invisible nucleon decay ($n \rightarrow \nu\nu\bar{\nu}$) mode.