

PROPOSAL FOR EXPERIMENT AT RCNP

March 11, 2002

Study of three-body force via the measurement of spin correlation coefficients $C_y^{y',y'}$ and $C^{y',y'}$ in $p+d$ elastic scattering.

SPOKESPERSON:

Name Atsushi Tamii
 Institution Department of Physics, University of Tokyo
 Title or Position Research Associate
 Address 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.
 Phone number +81-(0)3-5841-4236
 FAX number +81-(0)3-5841-7642
 E-mail tamii@phys.s.u-tokyo.ac.jp

EXPERIMENTAL GROUP:

Name	Institution	Title or Position
Hideyuki Sakai	Dep. of Physics, Univ. of Tokyo	(P)
Kentaro Yako	Dep. of Physics, Univ. of Tokyo	(D2)
Kimiko Sekiguchi	Dep. of Physics, Univ. of Tokyo	(D2)
Seitaro Sakoda	Dep. of Physics, Univ. of Tokyo	(D1)
Hiromitsu Kato	Dep. of Physics, Univ. of Tokyo	(M2)
Michio Hatano	Dep. of Physics, Univ. of Tokyo	(M2)
Yukie Maeda	Dep. of Physics, Univ. of Tokyo	(M2)
Takaaki Saito	Dep. of Physics, Univ. of Tokyo	(M1)
Nobuyuki Uchigashima	Dep. of Physics, Univ. of Tokyo	(M1)
Kichiji Hatanaka	RCNP, Osaka Univ.	(P)
Tomotsugu Wakasa	RCNP, Osaka Univ.	(RA)
Junichiro Kamiya	RCNP, Osaka Univ.	(D1)
Daisuke Hirooka	RCNP, Osaka Univ.	(M2)
Youhei Shimizu	RCNP, Osaka Univ.	(M1)
Yasuyuki Kitamura	RCNP, Osaka Univ.	(M1)
Hiroyuki Okamura	Dep. of Physics, Saitama Univ.	(AP)
Tomohiro Uesaka	Dep. of Physics, Saitama Univ.	(RA)
Kenji Suda	Dep. of Physics, Saitama Univ.	(D1)
Jun Nishikawa	Dep. of Physics, Saitama Univ.	(M2)

RUNNING TIME: Test running time for experiment 4 days
 Data runs 6 days

BEAM LINE: Ring : WS course

BEAM REQUIREMENTS: Type of particle polarized p
 Beam energy 400 MeV
 Beam intensity 500 nA

BUDGET: Experimental expenses 6,500,000 yen
 Travel plans 14 participants should be supported by RCNP

Study of three-body force via the measurement of spin correlation coefficients $C_y^{y',y'}$ and $C_{y'}^{y',y'}$ in $p+d$ elastic scattering.

SPOKESPERSON: Atsushi Tamii

SUMMARY OF THE PROPOSAL

Three-nucleon (3N) effect in 3N state is of particular interest in nuclear physics. The 3N state can be exactly described by Faddeev equation with nucleon-nucleon (NN) interaction. 3N force effect can, therefore, be extracted by comparing precise experimental results and rigorous theoretical calculations using modern NN interactions. It has been found that discrepancy between the experimental data of pd elastic cross sections at intermediate energies and Faddeev calculations can be well reproduced by incorporating 3N force in the Faddeev calculation. But there still remains discrepancies in spin observables. The fact may imply defects in the spin dependence of the 3N force. 3N force effect is expected to be larger at higher energies, but experimental data are scarce at proton energies above 200 MeV especially for various spin observables.

We propose to measure three-spin correlation coefficient $C_{y(p)}^{y'(p),y'(d)}$ and two-spin correlation coefficient $C_{y'}^{y'(p),y'(d)}$ in pd elastic scattering at $E_p = 400$ MeV together with other spin observables $A_{y(p)}$, $P_{y(p)}$, $P_{y(d)}$, $K_{y(p)}^{y'(p)}$ and $K_{y(p)}^{y'(d)}$. The data will be taken in an angular range where the observables are expected to be most sensitive to the 3N force effect. These data provide important information on the 3N force in 3N systems at high energy ($E_p=400$ MeV).