

## PROPOSAL FOR EXPERIMENT AT RCNP

28 Aug. 2000

**TITLE:**Search for excited  $\alpha$ -cluster states in  ${}^6\text{Li}$  via  ${}^6\text{Li}$  ( ${}^7\text{Li}, {}^7\text{Be}, d$ ) reaction**SPOKESPERSON:**

Full Name Hidetoshi Akimune  
 Institution Department of Physics, Konan University  
 Title or Position Lecturer  
 Address Okamoto 8-9-1, Higashinada, Kobe 658-8501, Japan  
 Phone number +81-78-435-2470  
 FAX number +81-78-435-2539  
 E-mail akimune@konan-u.ac.jp

**EXPERIMENTAL GROUP:**

Full Name	Institution	Title or Position
Tamio Yamagata	Dep. of Physics, Konan Univ.	(P)
Hiroaki Utsunomiya	Dep. of Physics, Konan Univ.	(P)
Hidetoshi Akimune	Dep. of Physics, Konan Univ.	(L)
Kaoru Yamasaki	Dep. of Physics, Konan Univ.	(D1)
Ayako Shiokawa	Dep. of Physics, Konan Univ.	(M1)
Shintaro Nakayama	Dep. of Physics, Univ. of Tokushima	(P)
Masaru Yosoi	RCNP, Osaka Univ.	(RA)
Yasushi Arimoto	JAERI (SPring-8)	(RF)
Masayoshi Tanaka	Kobe Tokiwa College	(P)

**RUNNING TIME:** Data runs 5 days (including test running time)**BEAM LINE:** Ring : WS course

**BEAM REQUIREMENTS:** Type of particle  ${}^7\text{Li}^{3+}$   
 Beam energy 65 MeV/A  
 Beam intensity  $\leq 3$  nA  
 Any other requirements Energy stability over  
 running time

**BUDGET:** Experimental expenses 200,000 yen

**TITLE:****Search for excited  $\alpha$ -cluster states in  ${}^6\text{Li}$  via  ${}^6\text{Li} ({}^7\text{Li}, {}^7\text{Be}, \text{d})$  reaction****SPOKESPERSON:** Hidetoshi Akimune**SUMMARY OF THE PROPOSAL**

The present work aims at the search for an excited states of an  $\alpha$ -cluster in the  ${}^6\text{Li}$  nucleus ( excited-cluster states, ECS), i.e. the giant dipole resonance (GDR) and spin-dipole resonance (SDR) of an  $\alpha$ -cluster, by observing their analogue states in the  ${}^6\text{He}$  nucleus excited via the  ${}^6\text{Li} ({}^7\text{Li}, {}^7\text{Be})$  reaction. In our previous experiment with this reaction, a such possible peak for ECS has been found in singles spectra at an excitation energy about 28 MeV in  ${}^6\text{He}$ , riding on the broad bumps of known GDR and SDR of the  ${}^6\text{Li}$  nucleus itself.

We suppose that a deuteron-cluster also presents as a spectator in ECS. In a decay of this excited nucleus the deuteron-cluster should be emitted, while analogue states of the GDR and SDR in  ${}^6\text{He}$  decay mainly via a neutron emission. Thus, observing the deuterons decaying from excited states of  ${}^6\text{He}$  in coincidence with  ${}^7\text{Be}$  particles, ECS should be enhanced and should be discriminated from other non-cluster excitation.