

PROPOSAL FOR EXPERIMENT AT RCNP

15 January 2002

TITLE:**Three Nucleon Force Effects in $n + d$ elastic scattering at 250 MeV****SPOKESPERSON:**

Name Yukie Maeda
 Institution Department of Physics, University of Tokyo
 Title or Position Student (D1)
 Address 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.
 Phone number +81-(0)3-5841-4238
 FAX number +81-(0)3-5841-7642
 E-mail yukie@nucl.phys.s.u-tokyo.ac.jp

EXPERIMENTAL GROUP:

Name	Institution	Title or Position
Hideyuki Sakai	Dep. of Physics, Univ. of Tokyo	(P)
Atsushi Tamii	Dep. of Physics, Univ. of Tokyo	(RA)
Kentaro Yako	Dep. of Physics, Univ. of Tokyo	(D3)
Kimiko Sekiguchi	Dep. of Physics, Univ. of Tokyo	(D3)
Michio Hatano	Dep. of Physics, Univ. of Tokyo	(D1)
Takaaki Saito	Dep. of Physics, Univ. of Tokyo	(M2)
Nobuyuki Uchigashima	Dep. of Physics, Univ. of Tokyo	(D1)
Tomoyuki Ishida	Dep. of Physics, Univ. of Tokyo	(M1)
Kichiji Hatanaka	RCNP, Osaka Univ.	(P)
Yasuhiro Sakemi	RCNP, Osaka Univ.	(AP)
Tomotsugu Wakasa	RCNP, Osaka Univ.	(RA)
Junichiro Kamiya	RCNP, Osaka Univ.	(D2)
Youhei Shimizu	RCNP, Osaka Univ.	(M2)
Yasuyuki Kitamura	RCNP, Osaka Univ.	(M2)
Hiroyuki Okamura	Dep. of Physics, Saitama Univ.	(AP)
Kenji Suda	Dep. of Physics, Saitama Univ.	(D2)
Hiroyuki Kamada	Dep. of Physics, Kyushu Institute of Technology	(AP)
M.B. Greenfield	International Christian Univ.	(P)

RUNNING TIME: Test running time for experiment 4 days
 Data runs 11 days

BEAM LINE: N0 (NTOF + NPOL2)
 WS course + LAS

BEAM REQUIREMENTS: Type of particle p
 Beam energy 250 MeV
 Beam intensity N0 : 50 nA (after 1/9 pulsing)
 WS : 500 nA
 Time resolution < 500 ps (FWHM)

BUDGET:

Experimental expenses	5,000,000 yen
Travel plan	500,000 yen
Total	5,500,000 yen

TITLE:

Three Nucleon Force Effects in $n + d$ elastic scattering at 250 MeV

SPOKESPERSON: Yukie Maeda

SUMMARY OF THE PROPOSAL

Three-nucleon force (3NF) effect is one of the hot topics in nuclear physics. Theoretically, we can treat the three nucleon state exactly by solving Faddeev equation with modern nucleon-nucleon (NN) interaction. Therefore 3NF can be estimated by comparing precise experimental results and rigorous Faddeev calculations. Recently it has been found that the differential cross sections of elastic $p + d$ scattering at intermediate energy region can be well reproduced by incorporating 3NF in the Faddeev calculation. Furthermore, precise measurements of not only cross sections but also spin observables have been carrying out in $p + d$ scattering.

However, 3NF effects have been studied by comparing the $p + d$ experimental data with $n + d$ calculations because the inclusion of the Coulomb interaction into the calculation is very difficult. To study 3NF effects in a Coulomb-free system, we have performed the $\vec{n} + d$ elastic measurement at $E_n = 250$ MeV in the backward angular region $\theta_{cm} = 85^\circ - 180^\circ$.

We propose to measure the differential cross sections in $n + d$ elastic scattering at $E_n = 250$ MeV in the forward angular region. The data will be taken in an angular range where the observables are expected to be sensitive more directly to Coulomb force. These data together with the earlier one cover a full angular range and allow us to make a Coulomb free comparison to the Faddeev calculations for the first time at intermediate energy region. In addition to that, we can make a direct data-to-data comparison between this $n + d$ and existing