

PROPOSAL FOR EXPERIMENT AT RCNP

14 Jan. 2002

TITLE : Observation of ($p, 2p$) Reactions on Light Nuclei

SPOKESPERSONS : NORO, Tetsuo Professor
 Dept. of Phys., Kyushu University
 6-10-1 Hakozaki, Higashi, Fukuoka 812-8581
 Phone: 092-642-2544, Fax: 642-2553
 E-mail: noro@nucl.phys.kyushu-u.ac.jp

NAKAMURA, Masanobu Lecturer
 Dept. of Phys., Kyoto University
 Phone: 075-753-3853, Fax: 753-3887
 E-mail: nakamura@nh.scphys.kyoto-u.ac.jp

EXPERIMENTAL GROUP :

Sagara, K	Kyusyu U.	AP	Ishida, K	Kyushu U.	M2
Nozoe, S.	Kyushu U.	M1			
Hatanaka, K.	RCNP	P	Sakemi, Y.	RCNP	AP
Wakasa, T.	RCNP	RA	Yoshimura, M.	RCNP	Research Fellow
Yoshida, H. P.	RCNP	JSPS Fellow	Kamiya, J.	RCNP	D2
Shimizu, Y.	RCNP	M2	Fujita, K.	RCNP	M1
Sakamoto, N.	RCNP	M1			
Sakaguchi, H.	Kyoto U.	AP	Takeda, H.	Kyoto U.	D3
Yasuda, Y.	Kyoto U.	D1	Terashima, S.	Kyoto U.	M1

RUNNING TIME : Test running time and calibration runs 2.0 days
 Data runs 5.5 days

BEAM LINE : WS course (Grand Raiden + LAS)

BEAM REQUIREMENTS : Type of particle Polarized proton
 Beam energy 392 MeV
 Beam intensity 300 nA

BUDGET : Experimental expenses 7.3 M Yen

TITLE : Observation of $(p, 2p)$ Reactions on Light Nuclei

SPOKESPERSON : NORO, Tetsuo and NAKAMURA, Masanobu

SUMMARY OF THE PROPOSAL

It is proposed to measure differential cross sections and analyzing powers for $(p, 2p)$ reactions on nuclei from deuteron to p -shell or light sd -shell nuclei. One of the purposes of this measurement is to study the A_y reduction problem observed in this reaction corresponds to $s_{1/2}$ -knockout. From the proposed systematic measurement using light nuclei, it is expected to see whether the reduction of the analyzing powers of these reactions scales to the nuclear density or to the deviation from the on-shell condition caused by finite Q -values of the reactions. Helium-4 target, which central density is unusually high but the separation energy of one nucleon does not stand out from those of $1s_{1/2}$ nucleon in neighboring nuclei, will play an important role in this systematic study. Another purpose is to examine the reliability of IA analysis, where the recoil effect is neglected or only perturbedly treated, for such light target nuclei. For this purpose, measurement with various kinematical condition is planned for a few target.