

PROPOSAL FOR EXPERIMENT AT RCNP

22 July 2003

TITLE:Measurement of the α -decay from the states around $E_x=10.3$ MeV in ^{12}C **SPOKESPERSON:**

Full Name Masatoshi Itoh
 Institution Research Center for Nuclear Physics, Osaka University
 Title or Position Research Fellow
 Address 10-1, Mihogaoka Ibaraki, Osaka 567-0047, Japan
 Phone number +81-6-6879-8939
 FAX number +81-6-6879-8924
 E-mail itoh@rcnp.osaka-u.ac.jp

EXPERIMENTAL GROUP:

Full Name	Institution	Title or Position
M. Fujiwara	Research Center for Nuclear Physics	(AP)
M. Uchida	Research Center for Nuclear Physics	(PD)
K. Nakanishi	Research Center for Nuclear Physics	(S)
K. Kawase	Research Center for Nuclear Physics	(S)
H. Hashimoto	Research Center for Nuclear Physics	(S)
S. Okumura	Research Center for Nuclear Physics	(S)
H. Sakaguchi	Department of Physics, Kyoto University	(AP)
M. Yosoi	Department of Physics, Kyoto University	(A)
Y. Yasuda	Department of Physics, Kyoto University	(S)
S. Terashima	Department of Physics, Kyoto University	(S)
S. Kishi	Department of Physics, Kyoto University	(S)
J. Zenihiro	Department of Physics, Kyoto University	(S)
Y. Nakatsugawa	Department of Physics, Kyoto University	(S)
H. Akimune	Department of Physics, Konan, University	(AP)
T. Kawabata	CNS, University of Tokyo	(A)
U. Garg	University of Notre Dame	(P)
R.B.K. Nayak	University of Notre Dame	(PD)

RUNNING TIME: Installation time without beam 3 days
 Development of device 0 days
 Test running time for experiment 1 days
 Data runs 7 days

BEAM LINE:

Ring : WS course

BEAM REQUIREMENTS:

Type of particle α
 Beam energy 400 MeV
 Beam intensity ≤ 5 nA
 Any other requirements

energy resolution ≤ 200 keV

halo-free, stable beam position, small emittance

BUDGET:

Experimental expenses 2,000,000 yen

TITLE:**Measurement of the α -decay from states around $E_x=10.3$ MeV in ^{12}C** **SPOKESPERSON:** Masatoshi Itoh**SUMMARY OF THE PROPOSAL**

The ^{12}C nucleus is one of the most widely investigated nuclei in the entire nuclear chart. However, there are still many unanswered questions. Among them, questionable the multipolarity of the broad level at $E_x=10.3$ MeV in ^{12}C . This state has been tentatively assigned to be 0^+ . According to the 3- α RGM calculation by Kamimura, a 2^+ level should co-exist in this excitation energy region which should be a 2^+ member of the β band beginning with the 7.654 MeV 0^+ level in ^{12}C as the band head. These states has been predicted to be the molecule-like states consisting three α -particles.

In the E133 experiment, we found that the 2^+ state actually exists around $E_x=10.3$ MeV by the multipole decomposition analysis. However, we did not well reproduce the angular distribution obtained for the α -cluster 0^+ state at $E_x=7.654$ MeV. Therefore, the ambiguities still remain to distinguish the 2^+ state from the 0^+ state.

We wish to measure the decay α -particles from the state at $E_x=10.3$ MeV for ^{12}C to solve the long-standing question concerning the multipolarity. Decay studies add an important, and highly desirable dimension to investigation of the structure of the α -cluster state in that they provide information on their microscopic structure which is not obtainable from “inclusive” inelastic scattering experiments alone.