

PROPOSAL FOR EXPERIMENT AT RCNP

April 5, 2005

TITLE: Dipole resonances in ^4He **SPOKESPERSON:**

Shintaro NAKAYAMA (Professor of Physics)
 Department of Physics, University of Tokushima, Tokushima 770-8502
 e-mail: nakayama@ias.tokushima-u.ac.jp TEL: 088-656-7236

EXPERIMENTAL GROUP:

Shintaro NAKAYAMA	Dept. of Physics, Univ. of Tokushima	P
Kenichi FUSHIMI	Dept. of Physics, Univ. of Tokushima	AP
Rena HAYAMI	Dept. of Physics, Univ. of Tokushima	M1
Minoru SAKAMA	Rad. Sci. and Eng., Univ. of Tokushima	RA
Tamio YAMAGATA	Dept. of Physics, Konan University	P
Hidetoshi AKIMUNE	Dept. of Physics, Konan University	AP
Nobuyuki WARASHINA	Dept. of Physics, Konan University	M1
Masayoshi TANAKA	Kobe Tokiwa Jr. College	P
Mamoru FUJIWARA	RCNP, Osaka University	AP
Kousuke NAKANISHI	RCNP, Osaka University	D2
Keigo KAWASE	RCNP, Osaka University	D2
Hisanobu HASHIMOTO	RCNP, Osaka University	D1
Masaru YOSOI	Dept. of Physics, Kyoto University	RA
M.B. GREENFIELD	Dept. of Physics, ICU	P

RUNNING TIME:

Total running time not including beam preparation 6 days

BEAM LINE:

WS-course, Grand RAIDEN, NYMPHS

BEAM REQUIREMENTS:

Type of particle	$^7\text{Li}^{3+}$
Beam energy	455 MeV
Beam intensity	a few tens nA
Target	^4He , ^{12}C , Kapton foil
Other requirements	Energy resolution ~ 150 keV
Beam must be halo-free and stability over several days is required	

BUDGET:

Experimental expenses 600,000 yen
 Travel plans - 10 participants should be supported by RCNP

TITLE: Dipole resonances in ^4He

SPOKESPERSON:

Shintaro NAKAYAMA

Department of Physics, University of Tokushima, Tokushima 770-8502

SUMMARY OF THE PROPOSAL

We aim at investigating dipole resonances, i.e., the giant dipole resonance (GDR) and spin dipole resonance (SDR), in ^4He by using the ($^7\text{Li}, ^7\text{Be}$) reaction at 455 MeV on ^4He . The GDR and SDR of ^4He will be obtained by measuring the 0.43-MeV γ -ray of ^7Be in coincidence with ^7Be -scattered particles. These resonances are important for studies of the charge symmetry of nuclear force and the reaction mechanism.

The GDR in ^4He was investigated with the photonuclear reactions, (γ, n) and (γ, p), and widely discussed during the past few decades. However, reported observations of their cross sections and resonance shapes were in contradiction. Further there are scarce data for the SDR in ^4He . In the present experiment we will measure the excitation energies, resonance shapes, and widths for the GDR and SDR in ^4He .

The noble gas, ^4He , should be prepared as a gas target which is to be installed inside a cell with a Kapton foil window. ^7Li induced reactions on ^{12}C and the Kapton foil will be measured in order to account for the background they produce.