

## PROPOSAL FOR EXPERIMENT AT RCNP

21 July 2005

**TITLE:**

**Precise determination of Gamow-Teller  $\beta^-$  strengths on double beta decay nuclei**

**SPOKESPERSON:**

Masaki Sasano  
 School of Science The University of Tokyo  
 D1  
 Hongo 7-3-1, Bunkyo, Tokyo 113-0033, Japan  
 Phone number +81-3-5841-4234  
 FAX number +81-3-5841-7642  
 E-mail sasano@nucl.phys.s.u-tokyo.ac.jp

**EXPERIMENTAL GROUP:**

Full Name	Institution	Title or Position
Hideyuki Sakai	Department of Physics, The University of Tokyo	(P)
Kentaro Yako	Department of Physics, The University of Tokyo	(RA)
Hironori Kuboki	Department of Physics, The University of Tokyo	(D2)
Yoshiyuki Takahashi	Department of Physics, The University of Tokyo	(M2)
Tomotsugu Wakasa	Department of Physics, Kyushu University	(AP)
Masanori Dozono	Department of Physics, Kyushu University	(M1)
Kimiko Sekiguchi	RIKEN	(PD)
Tomohiro Uesaka	CNS, The University of Tokyo	(AP)
Takahiro Kawabata	CNS, The University of Tokyo	(RA)
Yukie Maeda	CNS, The University of Tokyo	(PD)
Satoshi Sakaguchi	CNS, The University of Tokyo	(M2)
Yoshiko Sasamoto	CNS, The University of Tokyo	(M2)
Kichiji Hatanaka	RCNP, Osaka University	(P)
Atsushi Tamii	RCNP, Osaka University	(AP)
Kazuo Muto	Department of Physics, Tokyo Institute of Technology	(AP)
Hiroyuki Okamura	CYRIC, Tohoku University	(P)
Dieter Frekers	Institute of Nuclear Physics, University Muenster	(P)
M.B Greenfield	The Division of Natural Science, International Christian University	(P)

**RUNNING TIME:** Setup runs 1 day  
 Data runs 15 days  
 Total 16 days

**BEAM LINE:** Ring : N0 course

**BEAM REQUIREMENTS:** Type of particle protons  
 Beam energy 300 MeV  
 Beam intensity  $\geq 100$  nA after the pulsing of  
 1/5  
 Any other requirements time width  $\leq 250$  ps  
 halo-free, small emittance

**BUDGET:**

Experimental expenses	100,000 yen
Travel plan	600,000 yen
Total	700,000 yen

**TITLE:**

**Precise determination of Gamow-Teller  $\beta^-$  strengths on double beta decay nuclei**

**SPOKESPERSON:** Masaki Sasano

### SUMMARY OF THE PROPOSAL

We propose to measure the  $(p, n)$  reactions at 300 MeV on  $^{48}\text{Ca}$ ,  $^{76}\text{Ge}$ ,  $^{100}\text{Mo}$  and  $^{116}\text{Cd}$  with the new neutron detection system NPOL3 at the neutron time-of-flight (NTOF) facility to extract the Gamow-Teller (GT)  $\beta^-$  strengths in a wide excitation energy region up to 50 MeV by using a multipole decomposition analysis (MDA).

The purpose of this proposal is two folds. One is to obtain the GT strengths by the  $(p, n)$  reaction from parent to intermediate daughter nuclei of the double beta decay for understanding the intermediate structure of the daughter nuclei. The  $(n, p)$  measurement from the grandchild to daughter nuclei will be followed in future. These data will be used to deduce the nuclear matrix elements of the  $2\nu\beta\beta$  decay.

Another is to test the proportionality relation between a  $B(\text{GT})$  value and a differential cross section at  $0^\circ$  for the charge exchange reactions. Recently we found a large discrepancy in the deduced  $B(\text{GT})$  value for  $^{58}\text{Cu}(E_x = 1.051 \text{ MeV})$  between the  $(^3\text{He}, t)$  ( $B(\text{GT}) = 0.265 \pm 0.013$ ) and the  $(p, n)$  ( $B(\text{GT}) = 0.414 \pm 0.006$ ) reactions. It has been pointed out that the  $B(\text{GT})$  value for the  $^{116}\text{In}(gnd)$  state deduced by the  $(^3\text{He}, t)$  reaction,  $0.032 \pm 0.005$ , is largely different from that obtained from a  $\beta$ -decay experiment by a factor of 15. This discrepancy can be examined by the present proposed  $^{116}\text{Cd}(p, n)$  measurement and we can see whether the discrepancy of  $^{58}\text{Cu}(E_x = 1.051 \text{ MeV})$  is just a peculiar case or not.