

PROPOSAL FOR EXPERIMENT AT RCNP

January 20, 2005

TITLE:**Inelastic α scattering exciting the superdeformed band in ^{40}Ca and ^{32}S .****SPOKESPERSON:**

Full Name	Masatoshi ITOH
Cyclotron and Radioisotope Center, Tohoku University	
Title or Position	Research Associate
Address	6-3, Aramaki-Aza-Aoba, Aoba, Sendai, Miyagi 980-8578
Phone number	+81-22-795-7796
FAX number	+81-22-795-7997
E-mail	itoh@cyric.tohoku.ac.jp

EXPERIMENTAL GROUP:

Name	Institution	Title or Position
H. Sakaguchi	Department of Physics, Kyoto University	(AP)
T. Murakami	Department of Physics, Kyoto University	(RA)
Y. Yasuda	Department of Physics, Kyoto University	(D3)
Y. Terashima	Department of Physics, Kyoto University	(D3)
J. Zenihiro	Department of Physics, Kyoto University	(D1)
Y. Nakatsugawa	Department of Physics, Kyoto University	(D1)
Y. Iwao	Department of Physics, Kyoto University	(M2)
T. Emori	Department of Physics, Kyoto University	(M2)
M. Fujiwara	RCNP, Osaka University	(AP)
M. Yosoi	RCNP, Osaka University	(AP)
Y. Nakanishi	RCNP, Osaka University	(D3)
K. Kawase	RCNP, Osaka University	(D3)
H. Hashimoto	RCNP Osaka University	(D2)
U. Garg	Department of Physics, University of Notre Dame	(P)
M. Uchida	Department of Physics, Tokyo Institute of Technology	(RA)
H. Akimune	Department of Physics, Konan University	(AP)
T. Kawabata	CNS, University of Tokyo	(RA)

RUNNING TIME: Installation time without beam 1.5 days(for each beam time)
 Beam tuning and data runs 4 days

BEAM LINE: Ring : WS course

BEAM REQUIREMENTS: Type of particle alpha
 Beam energy 400MeV for α
 Beam intensity 20 nA
 Any other requirements energy resolution ≤ 150
 keV for α and ≤ 50 keV for high resolution α
 halo-free, small emittance

BUDGET: Experimental expenses 540,000 yen

TITLE:**Inelastic α scattering exciting the superdeformed band in ^{40}Ca and ^{32}S** **SPOKESPERSON:** M. Itoh**SUMMARY OF THE PROPOSAL**

During the beam time for ISGDR for light nuclei we have found for the first time candidate states of the superdeformed band in ^{32}S , which have been searched during these 20 years by many researchers in the world.

In this proposal, in order to study the transition form factor of the superdeformed band, we propose to measure inelastic α scattering on ^{40}Ca , which is already established by the measurement of the sequential γ decay from high spin state. At the next stage, we plan to measure inelastic α scattering on ^{32}S with the high resolution α beam exciting the same states to obtain the information on widths of these states by using the dispersion matching method. Our new data will also give new information on the origin of the superdeformed band, namely molecular like structures of ^{16}O - ^{16}O states in ^{32}S .