

PROPOSAL FOR EXPERIMENT AT RCNP

06/02/2007

TITLE: Systematic study of (p, n) reactions on light nuclei at 350 MeV**SPOKESPERSON:**

Tomotsugu WAKASA,
 Associate Professor,
 Department of Physics, Kyushu University,
 Hakozaki 6-10-1, Higashi, Fukuoka 812-8581, Japan
 Phone number : +81-92-642-2543
 FAX number : +81-92-642-2553
 E-mail : wakasa@phys.kyushu-u.ac.jp

EXPERIMENTAL GROUP:

| | | | | | |
|-----------------|--------------|----|--------------|--------------|---|
| <u>E. Ihara</u> | Kyushu Univ. | M | T. Noro | Kyushu Univ. | P |
| M. Dozono | Kyushu Univ. | M | Y. Yamada | Kyushu Univ. | M |
| H. Tanabe | Kyushu Univ. | M | K. Sagara | Kyushu Univ. | P |
| S. Kuroita | Kyushu Univ. | M | T. Sugimoto | Kyushu Univ. | M |
| K. Hatanaka | RCNP | P | H. Okamura | RCNP | P |
| A. Tamii | RCNP | AP | Y. Tameshige | RCNP | D |
| H. Matsubara | RCNP | D | M. Kato | RCNP | M |
| Y. Sakemi | CYRIC | P | K. Sekiguchi | RIKEN | R |

RUNNING TIME:

| | |
|-------------------------------------|-----------|
| Development of (p, n) measurement | 3.0 days |
| Calibration of NPO,L3 | 1.0 day |
| Measurement of σ and A_y | 4.0 days |
| Measurement of elastic scattering | 1.0 day |
| Measurement of $D_{ii}(0^\circ)$ | 4.0 days |
| Total | 13.0 days |

BEAM LINE: WS and N0 (WS + GR + NPOL3/N0 + NPOL3)**BEAM REQUIREMENTS:**

| | |
|-------------------|---|
| Type of particle | Polarized Protons |
| Beam energy | $\simeq 350$ MeV |
| Beam intensity | > 500 nA on target before pulse selection |
| Time resolution | < 300 ps (FWHM) |
| Beam polarization | > 0.7 |
| Injection Mode | High Current Mode |
| Pulse selection | 1/9 or 1/2 |

BUDGET:

| | |
|---------------------------|-----------|
| Summary of budget request | 4,700,000 |
| Experimental expenses | 3,900,000 |
| Travel plan | 800,000 |

TITLE: Systematic study of (p, n) reactions on light nuclei at 350 MeV
SPOKESPERSON: Tomotsugu WAKASA

SUMMARY OF THE PROPOSAL

We propose to measure (p, n) reactions on ${}^3\text{He}$ and ${}^4\text{He}$ at $T_p \simeq 350$ MeV. The data with the existing ${}^2\text{H}(p, n)$ data provide the systematic data for (p, n) reactions on light nuclei of $A=2-4$ at $T_p \simeq 350$ MeV where the reaction mechanism is expected to be simple and a distorted wave impulse approximation (DWIA) is reliable.

For ${}^4\text{He}$, an unexpected enhancement of the cross section has been observed in the quasielastic (QE) scattering at $T_p=197$ MeV. This enhancement might be a signature of high density ($>$ normal density) effects of ${}^4\text{He}$ which are not included in present standard theoretical calculations such as DWIA calculations employing random phase approximation (RPA) response functions and a density-dependent effective nucleon mass. One of these effects is the reduction of hadron mass and coupling constant with nuclear density as a signature of partial restoration of chiral symmetry in nuclear matter. However, there are ambiguities in theoretical calculations and their interpretations since the reaction mechanism might be complicated at high energy transfers ($T_n \lesssim 100$ MeV) of ${}^4\text{He}(p, n)$ at $T_p=197$ MeV. The reaction mechanism should be more simple at $T_p \simeq 350$ MeV. Thus the data at $T_p \simeq 350$ MeV will provide clear and useful information on high density effects in nuclear matter, and a characterization of these effects is expected to be crucial for understanding relativistic heavy ion collisions and neutron stars.

For ${}^3\text{He}$, our data will provide the information on both the $T = 3/2$ resonance in the three-nucleon system and the $T = 3/2$ three-nucleon forces. We can also refine the understanding for the structure and reactions of light nuclei (e.g. QE peak shift) because systematic data for (p, n) reactions on ${}^2\text{H}$ (RCNP-E59 and E131), ${}^3\text{He}$ and ${}^4\text{He}$ (this proposal) at the same incident energy of $T_p \simeq 350$ MeV and the reaction angles become available.