

**PROPOSAL FOR EXPERIMENT AT RCNP**

14 February 2008

**TITLE:****Search for Direct Evidence of Tensor Interaction:  
High Momentum Component in Nuclei****SPOKESPERSON:**

Full Name Isao TANIHATA  
 Institution Cosmonuclear Physics Division, RCNP, Osaka University  
 Title or Position Professor  
 Address 10-1 Mihogaoka, Ibaraki-shi, Osaka 567-0047, Japan.  
 Phone number +81-6-6879-8918  
 FAX number +81-6-6879-8899  
 E-mail tanihata@rcnp.osaka-u.ac.jp

Full Name Hooi Jin ONG  
 Institution Cosmonuclear Physics Division, RCNP, Osaka University  
 Title or Position Assistant Professor  
 Address 10-1 Mihogaoka, Ibaraki-shi, Osaka 567-0047, Japan.  
 Phone number +81-6-6879-8919  
 FAX number +81-6-6879-8899  
 E-mail onghjin@rcnp.osaka-u.ac.jp

**EXPERIMENTAL GROUP:**

Name	Institution	Title or Position
H. SAKAGUCHI	Dep. of Applied Physics, Miyazaki Univ.	Professor
K. MATSUTA	Dep. of Physics, Osaka Univ.	Associate Professor
M. FUKUDA	Dep. of Physics, Osaka Univ.	Associate Professor
M. MIHARA	Dep. of Physics, Osaka Univ.	Assistant Professor
D. NISHIMURA	Dep. of Physics, Osaka Univ.	M2
A. OZAWA	Tsukuba Univ.	Associate Professor
K. SEKIGUCHI	RIKEN Nishina Center	Researcher
H. OKAMURA	RCNP	Professor
A. TAMII	RCNP	Associate Professor
M. YOSOI	RCNP	Associate Professor
K. SUDA	RCNP	Researcher
T. ADACHI	RCNP	Researcher
Y. TAMESHIGE	RCNP	D3
H. MATSUBARA	RCNP	D2
D. ISHIKAWA	RCNP	M1

**THEORETICAL GROUP:**

Name	Institution	Title or Position
K. IKEDA	RIKEN Nishina Center	Emeritus Professor
H. TOKI	RCNP	Professor
T. MYO	RCNP	Researcher
Y. OGAWA	RCNP	Researcher

**RUNNING TIME:**

GR set up and tuning + change of target + contingency 1.5+0.5+1.0 days  
(p,d) reaction runs for three targets 3.0 days  
(p,pd) reaction runs for two targets 4.0 days

**BEAM LINE:** Ring : WS beam line and Grand Raiden Spectrometer.

**BEAM REQUIREMENTS:**

Type of particle p  
Beam energy 200 MeV, 300 MeV, 392 MeV  
Beam intensity 10 nA

**OTHER REQUIREMENTS:**

Achromatic beam with resolution  $\leq 80$  keV  
Single turn and halo-free beam with small emittance

**BUDGET:**

Assembly of proton detector system 3,000 kyen  
Replacement of sliding membrane of the scattering chamber 300 kyen  
Travelling expenses including accommodation of 7 participants are to be provided by RCNP

**TITLE:**

**Search for Direct Evidence of Tensor Interaction:  
High Momentum Component in Nuclei**

**SPOKESPERSON:** Isao TANIHATA, Hooi Jin ONG

### SUMMARY OF THE PROPOSAL

Measurements of (p,d) and (p,pd) reactions on  $^{12}\text{C}$  and  $^{16}\text{O}$  targets using proton beams at 200 MeV, 300 MeV and 392 MeV are proposed in search of direct evidence of tensor interaction in high-momentum component in nuclei. The experiment will be performed using the Grand Raiden spectrometer at several finite angles between  $1^\circ$  and  $30^\circ$ . The main objective is to deduce the internal momentum distributions, specifically those of the s-wave neutrons in  $^{12}\text{C}$  and  $^{16}\text{O}$ , around  $2\text{ fm}^{-1}$  where the effects of tensor force are expected to be different. For this purpose, excitation energy up to 4 MeV, with resolution about 100 keV sufficient to separate most of the low lying states, will be measured to cover the s-hole states in  $^{11}\text{C}$  and  $^{15}\text{O}$ . Calibration will be carried out by measuring the (p,d) reaction on deuteron target at similar beam energies and angular settings.

These measurements will be performed with natural carbon foils, thin ice  $\text{H}_2\text{O}$  targets and a thin foil  $\text{CD}_2$  target. The proton beam will be stopped at a Faraday cup placed downstream of the Q1 magnet for the measurements at  $1^\circ$  and another Faraday cup placed inside the scattering chamber for the measurements at other angles. The standard focal plane detector system with two multi-wire drift chambers and two 10 mm-thick plastic scintillators will be used. For the (p,pd) reaction, a new  $\Delta E$ - $E$  detector system for the recoil protons will be installed in the backward angles with respect to the beam axis.

The cross sections for the (p,d) reaction at the measuring angles are expected to be about  $20\ \mu\text{b}/\text{sr}$  to  $100\ \mu\text{b}/\text{sr}$ . Assuming the cross section to be  $20\ \mu\text{b}/\text{sr}$ , we estimated the time required to obtain about 2000 counts of deuteron events.