

PROPOSAL FOR EXPERIMENT AT RCNP

13 July 2009

TITLE:**Stellar neutron sources and the s-process in massive stars.**

Spokespersons: G.P.A. Berg

University of Notre Dame, Notre Dame, IN 46556, USA

Ph: 001-219-631-6238, FAX: 001-219-631-5952, E-Mail: gpberg@bergs.edu

M. Wiescher,

University of Notre Dame, Notre Dame, IN 46556, USA

Ph: 001-219-631-6788, FAX: 001-219-631-5952, E-Mail: wiescher.1@nd.edu

Experimental Group: T. Adachi, Research Associate, KVI, Groningen, The Netherlands

G.P.A. Berg, Research Professor, Univ. of Notre Dame, USA

M. Couder, Research Associate, Univ. of Notre Dame, USA

Y. Fujita, Associate Professor, Dept. of Physics, Osaka Univ., Japan

H. Fujita, Researcher, RCNP, Osaka Univ., Japan

J. Görres, Research Professor, Univ. of Notre Dame, USA

M.N. Harakeh, Professor, KVI, Groningen, The Netherlands

K. Hatanaka, Professor, RCNP, Osaka Univ., Japan

S. O'Brien, Grad. Student, Univ. of Notre Dame, USA

Ong Hooi Jin, Assistant Professor, RCNP, Osaka Univ., Japan

A. Matic, Research Associate, FZD, Dresden, Germany

H. Matsubara, D3, RCNP, Osaka Univ., Japan

H. Okamura, Professor, RCNP, Osaka Univ., Japan

H. Schatz, Professor, Michigan State Univ., USA

Y. Sakemi, Professor, CYRIC, Tohoku University Japan

Y. Shimbara, Assistant Professor, Niigata University, Japan

Y. Shimizu, Researcher, CNS, University of Tokyo, Japan

A. Tamii, Associate Professor, RCNP, Osaka Univ., Japan

T. Wakasa, Associate Professor, Kyushu Univ., Japan

M. Wiescher, Professor, Univ. of Notre Dame, USA

M. Yosoi, Associate Professor, RCNP, Osaka Univ., Japan

Running Time: A total of 6 days running time is requested of which 2 days are needed for cyclotron optimization (momentum spread) beam line matching, spectrometer setup, and 4 days for production and calibration measurements including the time needed for mode changes (0° mode, Faraday cups behind Q1 and in scattering chamber) and gas target changes (^{18}O and ^{22}Ne gas and solid target).

Beam Line: Dispersive WS beam line and Grand Raiden Spectrometer in three modes (0° mode, Faraday cups behind Q1 and in scattering chamber).

Beam Requirements: Particle type and energy: ^4He of 206 MeV

Beam intensity and energy spread: A maximum of 100 pA is required, beam energy spread of the order of 100 keV or better.

Other requirements: Single turn halo-free beam, fully disp. matched beam on GR target.

Equipment required: Only existing and standard GR and focal plane equipment is needed

TITLE:

Stellar neutron sources and the s-process in massive stars.

SPOKESPERSON: G.P.A. Berg and M. Wiescher

SUMMARY OF THE PROPOSAL

• **Proposed experiment:** Measurements of (α, α') on ^{18}O and ^{22}Ne gas targets and a ^{26}Mg target foil with astrophysical motivation are proposed using the WS course and Grand Raiden (GR). GR will be used in three Faraday cup modes including the 0° mode with a Faraday cup downstream of the Focal plane detector.

This experiment is part of an astrophysics program at RCNP. It aims at resonance states above the α -threshold around 10 MeV excitation energy and is only possible with the high-resolution spectrometer since a resolution of 20 - 30 keV is required to resolve high-lying excited in the final nucleus. Two of the targets are gas targets and we will make use of the special, flat gas-target cell that allows dispersion matching as demonstrated previously. Targets will be of the order of 2 - 3 mg/cm² for best possible resolution.

• **Targets:** The targets of astrophysical interest are T = 1 nuclei in the sd-shell and we propose in this experiment measurements on ^{18}O , ^{22}Ne and ^{26}Mg targets. The target thicknesses of about 2 - 3 mg/cm² are a compromise of count rate and good resolution requirements.

• **Apparatus and Beam Properties:** The WS course in dispersive mode and the Grand Raiden spectrometer with the standard VDC focal plane detector system will be used. A stack of 3 mm, 10 mm, and 6 mm thick ΔE plastic scintillator will provide energy loss and timing signals for particle identification in the first two detectors for α -particles. Deuterons and protons will pass through the first two detectors and provide a veto signal in the third, 6 mm thick detector.

• **Beam time request:** The total beam time request of 6 days will be used as follows:

a) 2 days for beam preparation, detector and particle identification verifications, ion-optical setup and dispersion matching.

b) 3 days for measurements on all three targets. Count rates are about 200 events per hour for a ^{26}Mg target, an estimated cross section of 1 $\mu\text{b}/\text{sr}$, a solid angle of 2.4 msr solid angle, a detector efficiency of 0.8 and a beam current of 100 pA.

c) 1 day is needed for measurements on calibration targets.