

PROPOSAL FOR EXPERIMENT AT RCNP

10 July 2009

TITLE: Investigation of the Giant Monopole Resonance in the Lead Isotopes: Testing the MEM Effect on Nuclear Incompressibility

SPOKESPERSONS:

U. Garg, Physics Department, University of Notre Dame, Notre Dame, IN 46556, USA

e-mail: garg@nd.edu

FAX: 1.574.631.5952

M. Fujiwara, Research Center for Nuclear Physics, Osaka University, Suita, Osaka 567-0047, Japan

e-mail: fujiwara@rcnp.osaka-u.ac.jp

FAX: (06)6879-8899

EXPERIMENTAL GROUPS:

University of Notre Dame, USA - Research Center for Nuclear Physics, Japan - Konan University, Japan - Tohoku University, Japan - Kyoto University, Japan - Tokyo Institute of Technology, Japan - IPN, Orsay, France - KVI, Groningen, The Netherlands.

RUNNING TIME:

Total running time not including beam preparation 6 days

BEAM LINE: Grand Raiden

BEAM REQUIREMENTS:

Type of particle ^4He

Beam energy 400 MeV

Beam intensity 2 nA–30 nA

Other requirements beam must be halo-free

highest stability over the running period is required

BUDGET:

Purchase of enriched targets 500,000 Yen for ^{204}Pb , ^{206}Pb , ^{208}Pb isotopes

TITLE:

Investigation of the Giant Monopole Resonance in the Lead Isotopes: Testing the MEM Effect on Nuclear Incompressibility

SPOKESPERSONS:

U. Garg, Physics Department, University of Notre Dame, Notre Dame, IN 46556, USA

M. Fujiwara, Research Center for Nuclear Physics, Osaka University, Suita, Osaka 567-0047, Japan

SUMMARY OF THE PROPOSAL

We request beam time to measure the Giant Monopole Resonance (GMR) in the Lead isotopes, ^{204}Pb , ^{206}Pb , and ^{208}Pb . The primary aim behind these measurements is to test an intriguing idea put forward recently regarding nuclear incompressibility in doubly-magic nuclei: In analogy with nuclear masses, it has been suggested that there would be an effect of mutually-enhanced magicity (MEM) in the K_A and, hence, in the energies of the GMR's in doubly-magic nuclei with respect to the nearby isotopes. We believe it is imperative (and urgent) to perform precise systematic measurements of the centroid energies of the GMR in ^{204}Pb , ^{206}Pb , and ^{208}Pb in order to test the MEM effect on nuclear incompressibility.