

PROPOSAL FOR EXPERIMENT AT RCNP

11 July 2009

TITLE:

Search for high-spin isomeric states in N=80 isotones

SPOKESPERSON:

Full Name Costel Petrache
 Institution Department of Physics, University Paris Sud and IPN Orsay, France
 Title or Position Professor
 Address 15, rue G.Clemenceau, 91406 Orsay, France
 Phone number +33-1-6915-5153
 FAX number +33-1-6915-6470
 E-mail petrache@ipno.in2p3.fr

EXPERIMENTAL GROUP:

Full Name	Institution	Title or Position
D. Beaumel, J.A. Scarpaci, N. De Sereville	IPN Orsay	Researcher
A. Matta	IPN Orsay	Graduate student
N. Redon, O. Stezowski	IPN Lyon	Researcher
D. Guinet, P. Lautesse	IPN Lyon	Professor
A. Vancraeynest	IPN Lyon	Graduate student
P. Desesquelles	CSNSM Orsay	Professor
A. Korichi	CSNSM Orsay	Researcher
D. Curien, G. Duchene	IPHC Strasbourg	Researcher
L. Gaudefroy	CEA, DAM, Arpajon	Researcher
T. Shimoda	Dep. of Physics, Osaka Univ.	Professor
A. Odahara	Dep. of Physics, Osaka Univ.	Associate Professor
T. Suzuki	RCNP, Osaka Univ.	Post-doc
K. Tajiri	Dep. of Physics, Osaka Univ.	D1
M. Kazato, M. Suga, A. Takashima	Dep. of Physics, Osaka Univ.	M2
Y. Ito, Y. Kenmoku, K. Yamaguchi	Dep. of Physics, Osaka Univ.	M1
E. Ideguchi	CNS, University of Tokyo	Associate Professor
S. Go	CNS, University of Tokyo	M1
T. Morikawa	Dep. of Physics, Kyushu Univ.	Assistant Professor

RUNNING TIME: Installation time without beam 7 days
 RI beam tuning 1 days
 Data runs 10 days
 In total, we request 11 days.

BEAM LINE: Ring : EN course

BEAM REQUIREMENTS: Type of particle ^{18}O
 Beam energy 66 MeV (9.2 MeV/u)
 Beam intensity 3 pμA

BUDGET: Experimental expenses 2,700,000 yen

TITLE:**Search for high-spin isomeric states in N=80 isotones****SPOKESPERSON:** Costel Petrache**SUMMARY OF THE PROPOSAL**

We propose to investigate high-spin metastable states in nuclei around the N=82 shell closure, which having simple configurations, constitute ideal laboratories for the study of excited states in mesoscopic quantum systems. The comparison of the experimental and theoretical excitation energy of the many-quasiparticle isomers will allow us to extract the amount of pairing correlations and their reduction with seniority in the highly excited states of the nucleonic superfluid system.

In the present experiment we want to study high-spin states in the N=80 nucleus ^{136}Ba , which being close to stability, can be populated at high spins using the unique fusion-evaporation reaction employing the neutron-rich ^{17}N radioactive beam available at RCNP and the heaviest Sn isotope ^{124}Sn as target. There is no other stable projectile-target combination which can populate high-spin states in ^{136}Ba .

The identification of the predicted 14+ and 22+ high-spin isomers in ^{136}Ba will proof the yrastness of the maximum aligned multi-quasiparticle configurations in the N = 80 nuclei, in which only the 20+ isomer in ^{140}Nd is presently known. From the comparison with theoretical predictions of schematic and self-consistent models we will deduce the strength of the pairing correlations and find out if they persist in high-spin isomers as observed above N = 82. The study of the ^{136}Ba nucleus with Z=56, having the Fermi surface in the middle of the d5/2/g7/2 sub-shell (between Z=50 and Z=64), is essential for the understanding of the evolution of the pairing correlations. In fact, the very different occupation of the d5/2/g7/2 sub-shell in the two nuclei (^{136}Ba has 8 proton holes in Z=64, while ^{140}Nd only 4), is expected to lead to sizable differences in the pairing correlations in the high-seniority maximum aligned states. The present experiment is also essential for our research program aimed to the identification of high-spin isomers in the N=80 series of isotopes, which will finally allow the extraction of the proton pairing gap Δ_p in high-seniority states from the odd-even mass differences. The ^{17}N RI-beam will be delivered at the EN course by using the $^9\text{Be}(^{18}\text{O},^{17}\text{N})^{10}\text{B}$ primary reaction. The separated ^{17}N secondary beam will be transported through the spectrometer at F2 to bombard a ^{124}Sn target of 10 mg/cm². The ^{136}Ba nuclei populated by the p4n reaction channel will be selected through the particle-coincidences between the Ge detectors and a charged particle detector array. The high-spin isomers will be searched for by using delayed-coincidences at the focal plane. Prompt-delayed-coincidences will also be registered and will be used to identify structures built above the isomeric state.

We request 1 day for the ^{17}N RI-beam preparation and 10 days of data acquisition.