

Two-meson cloud contribution to the baryon antidecuplet binding



Tetsuo Hyodo^a,


A. Hosaka^a, F. J. Llanes-Estrada^b, E. Oset^c,
J. R. Peláez^b and M. J. Vicente Vacas^c

RCNP, Osaka^a *Madrid*^b *IFIC, Valencia*^c 2004, Dec. 23rd₁

Contents

★ **Introduction and motivations**

★ **Effective Lagrangian**



★ **Self-energy of antidecuplet**


★ **Mass shifts**

★ **Decay widths**

★ **Reaction cross section**

★ **Summary and conclusions**

Introduction : Exotic activities



taken from K. Goeke, et al., hep-ph/0411195

Introduction : Flavor SU(3) symmetry

Flavor SU(3) symmetry and its breaking by the strange quark mass

-> **Phenomenologically well realized**

Gell-Man—Okubo mass formulae ~ 3%

**N(938), $\Lambda(1116)$, $\Sigma(1192)$, $\Xi(1320)$,
 $\Delta(1232)$, $\Sigma(1385)$, $\Xi(1530)$, $\Omega(1670)$**


Existence of Θ^+ + Flavor SU(3) symmetry

→ **Existence of flavor partners of Θ^+**

Introduction : determining the quantum numbers

We have proposed $K^+ p \rightarrow \pi^+ \Theta^+ \rightarrow \pi^+ K^+ n (K^0 p)$

Polarization observable
can be used to determine
the quantum numbers



T. H., A. Hosaka, E. Oset,
Phys. Lett. B579, 290 (2004)

→ Understanding of reaction mechanism

Motivations

**Results of $\pi^- p \rightarrow K^- \Theta^+$ reaction at KEK
Total cross section $\sim 2 \mu b$**

K. Miwa, talk given at PENTAQUARK04


Why so small? What about $K^+ p \rightarrow \pi^+ \Theta^+$?

Possibility of $\Theta^+ \sim K\pi N$ bound state

P. Bicudo, et al., Phys. Rev. C69, 011503 (2004)

T. Kishimoto, et al., hep-ex/0312003

F. J. Llanes-Estrada, et al., Phys. Rev. C69, 055203 (2004)




Two-meson coupling



$$\Theta^+ \rightarrow KN$$

Very narrow




$$\Theta^+ \rightarrow K\pi N$$

Forbidden

$$N(1710) \rightarrow \pi\pi N$$

40–90 %



Large??

**Effective interactions which account
for the $N(1710) \rightarrow \pi\pi N$ decay**

Criteria to construct the Lagrangian

Interaction is flavor SU(3) symmetric
Chiral symmetric? -> later

Small number of derivatives
low energy : OK

Assumptions for Θ^+

N(1710) is the S=0 partner of antidecuplet
-> $J^P = 1/2^+$


No mixing with 8, 27,...

T.D. Cohen, Phys. Rev. D70, 074023 (2004)

S. Pakvasa and M. Suzuki, Phys. Rev. D70, 036002 (2004)

S. Ceci, et al., nucl-th/0406055

SU(3) structure of effective Lagrangian



$$8_M \otimes 8_M \otimes 8_B = (1 \oplus 8^s \oplus 8^a \oplus 10 \oplus 1\bar{10} \oplus 27)_{MM} \otimes 8_B$$

$$= 8 \quad \leftarrow \text{from } 1_{MM} \otimes 8_B$$

$$\oplus (1 \oplus 8 \oplus 8 \oplus \textcircled{10} \oplus 1\bar{10} \oplus 27) \quad \leftarrow \text{from } \underline{8^s_{MM}} \otimes 8_B$$

$$\oplus (1 \oplus 8 \oplus 8 \oplus \textcircled{10} \oplus 1\bar{10} \oplus 27) \quad \leftarrow \text{from } \underline{8^a_{MM}} \otimes 8_B$$

$$\oplus (8 \oplus \textcircled{10} \oplus 27 \oplus 35) \quad \leftarrow \text{from } \underline{10_{MM}} \otimes 8_B$$

$$\oplus (8 \oplus 1\bar{10} \oplus 27 \oplus 35') \quad \leftarrow \text{from } \underline{1\bar{10}_{MM}} \otimes 8_B$$

$$\oplus (8 \oplus \textcircled{10} \oplus 1\bar{10} \oplus 27 \oplus 27 \oplus 35 \oplus 35'' \oplus 64) \quad \leftarrow \text{from } \underline{27_{MM}} \otimes 8_B$$

Interaction Lagrangians 1

Antidecuplet field

$$P^{333} = \sqrt{6}\Theta_{\bar{10}}^+$$

$$P^{133} = \sqrt{2}N_{\bar{10}}^0 \quad P^{233} = -\sqrt{2}N_{\bar{10}}^+$$

$$P^{113} = \sqrt{2}\Sigma_{\bar{10}}^- \quad P^{123} = -\Sigma_{\bar{10}}^0 \quad P^{223} = -\sqrt{2}\Sigma_{\bar{10}}^+$$

$$P^{111} = \sqrt{6}\Xi_{\bar{10}}^{--} \quad P^{112} = -\sqrt{2}\Xi_{\bar{10}}^- \quad P^{122} = \sqrt{2}\Xi_{\bar{10}}^0 \quad P^{222} = -\sqrt{6}\Xi_{\bar{10}}^+$$

Meson and baryon fields

$$\Phi = \begin{pmatrix} \frac{1}{\sqrt{2}}\pi^0 + \frac{1}{\sqrt{6}}\eta & \pi^+ & K^+ \\ \pi^- & -\frac{1}{\sqrt{2}}\pi^0 + \frac{1}{\sqrt{6}}\eta & K^0 \\ K^- & \bar{K}^0 & -\frac{2}{\sqrt{6}}\eta \end{pmatrix}$$

$$B = \begin{pmatrix} \frac{1}{\sqrt{2}}\Sigma^0 + \frac{1}{\sqrt{6}}\Lambda & \Sigma^+ & p \\ \Sigma^- & -\frac{1}{\sqrt{2}}\Sigma^0 + \frac{1}{\sqrt{6}}\Lambda & n \\ \Xi^- & \Xi^0 & -\frac{2}{\sqrt{6}}\Lambda \end{pmatrix}$$

Interaction Lagrangians 2

Construction of 8s Lagrangian

$$\begin{aligned} D_i{}^j [8_{MM}^s] &= \phi_i{}^a \phi_a{}^j + \phi_i{}^a \phi_a{}^j - \frac{2}{3} \delta_i{}^j \phi_a{}^b \phi_b{}^a \\ &= 2\phi_i{}^a \phi_a{}^j - \frac{2}{3} \delta_i{}^j \phi_a{}^b \phi_b{}^a \end{aligned}$$

$$T^{ijk} [\bar{\mathbf{10}}_{BMM(8s)}] = 2\phi_l{}^a \phi_a{}^i B_m{}^j \epsilon^{lmk} + (i, j, k \text{ symmetrized})$$

→ $\mathcal{L}^{8s} = \frac{g^{8s}}{2f} \bar{P}_{ijk} \epsilon^{lmk} \phi_l{}^a \phi_a{}^i B_m{}^j + h.c.$

Interaction Lagrangians 3

Terms without derivative

$$\mathcal{L}^{8s} = \frac{g^{8s}}{2f} \bar{P}_{ijk} \epsilon^{lmk} \phi_l^a \phi_a^i B_m^j + h.c.$$

8s

$$\mathcal{L}^{8a} = 0$$

$$\mathcal{L}^{10} = 0$$

<- symmetry under exchange of mesons

$$\mathcal{L}^{27} = \frac{g^{27}}{2f} \left[4\bar{P}_{ijk} \epsilon^{lbk} \phi_l^i \phi_a^j B_b^a - \frac{4}{5} \bar{P}_{ijk} \epsilon^{lbk} \phi_l^a \phi_a^j B_b^i \right] + h.c.$$

Experimental information

$$N(1710) \rightarrow \pi\pi (s\text{-wave}, I=0) N$$

$$N(1710) \rightarrow \pi\pi (p\text{-wave}, I=1) N$$

With one derivative

8a



$$\mathcal{L}^{8a} = i \frac{g^{8a}}{4f^2} \bar{P}_{ijk} \epsilon^{lmk} \gamma^\mu (\partial_\mu \phi_l^a \phi_a^i - \phi_l^a \partial_\mu \phi_a^i) B_m^j + h.c.$$

Diagrams for self-energy

Real part : mass shift

Imaginary part : decay width

SU(3) breaking: masses of particles





$N(1710) \rightarrow \pi\pi(s\text{-wave}, I = 0)N$ **25 MeV**

$N(1710) \rightarrow \pi\pi(p\text{-wave}, I = 1)N$ **15 MeV**

→ $g^{8s} = 1.88$, $g^{8a} = 0.315$

Results of self-energy : Real part (mass shift)



All mass shifts are attractive.

More bound for larger strangeness.

Mass difference between Ξ and Θ

$\rightarrow 60 \text{ MeV} : \sim 20\% \text{ of } 320 = 1860 - 1540$

Results of self-energy : Imaginary part (decay width)

Decay [MeV]	$\Gamma^{(8s)}$	$\Gamma^{(8a)}$	$\Gamma_{BMM}^{(tot)}$
$N(1710) \rightarrow N\pi\pi$ (inputs)	25	15	40
$N(1710) \rightarrow N\eta\pi$	0.58	-	
$\Sigma(1770) \rightarrow N\bar{K}\pi$	4.7	6.0	24
$\Sigma(1770) \rightarrow \Sigma\pi\pi$	10	0.62	
$\Sigma(1770) \rightarrow \Lambda\pi\pi$	-	2.9	
$\Xi(1860) \rightarrow \Sigma\bar{K}\pi$	0.57	0.46	2.1
$\Xi(1860) \rightarrow \Xi\pi\pi$	-	1.1	

Other possible Lagrangians

$$\mathcal{L}^{8s} = \frac{g^{8s}}{2f} \bar{P}_{ijk} \epsilon^{lmk} \phi_l{}^a \phi_a{}^i B_m{}^j + h.c.$$

Two-meson 27 interaction

$$\mathcal{L}^{27} = \frac{g^{27}}{2f} \left[4 \bar{P}_{ijk} \epsilon^{lbk} \phi_l{}^i \phi_a{}^j B_b{}^a - \frac{4}{5} \bar{P}_{ijk} \epsilon^{lbk} \phi_l{}^a \phi_a{}^j B_b{}^i \right] + h.c.$$

Chiral symmetric interaction

$$\mathcal{L}^\chi = \frac{g^\chi}{2f} \bar{P}_{ijk} \epsilon^{lmk} (A_\mu)_l{}^a (A^\mu)_a{}^i B_m{}^j + h.c.$$

$$A_\mu = \frac{i}{2} (\xi^\dagger \partial_\mu \xi - \xi \partial_\mu \xi^\dagger) = -\frac{\partial_\mu \phi}{\sqrt{2}f} + \mathcal{O}(p^3) \quad \xi = e^{i\phi/\sqrt{2}f}$$

$$(A_\mu)_l{}^a (A^\mu)_a{}^i \rightarrow \frac{1}{2f^2} \partial_\mu \phi_l{}^a \partial^\mu \phi_a{}^i$$

SU(3) breaking interaction $M = \text{diag}(\hat{m}, \hat{m}, m_s)$

$$\mathcal{L}^M = \frac{g^M}{2f} \bar{P}_{ijk} \epsilon^{lmk} S_l{}^i B_m{}^j$$

$$S = \xi M \xi + \xi^\dagger M \xi^\dagger = \mathcal{O}(\phi^0) - \frac{1}{2f^2} (2\phi M \phi + \phi \phi M + M \phi \phi) + \mathcal{O}(\phi^4)$$

Chiral symmetric Lagrangian

$$\mathcal{L}^{8s} = \frac{g^{8s}}{2f} \bar{P}_{ijk} \epsilon^{lmk} \phi_l{}^a \phi_a{}^i B_m{}^j + h.c.$$

$$\mathcal{L}^{\chi(2)} = \frac{g^\chi}{2f} \bar{P}_{ijk} \epsilon^{lmk} \frac{1}{2f^2} \partial_\mu \phi_l{}^a \partial^\mu \phi_a{}^i B_m{}^j + h.c.$$

SU(3) structure : Identical !

**Only loop integral is changed
<- adjusting the cutoff, we would have
the same results**

N(1710) decay -> $g^\chi = 0.218$

Results of chiral Lagrangian

[MeV]

$\text{Re}\{\Sigma\}$	8s	$\chi(2)$
Θ	-100	-99
N	-87	-83
Σ	-79	-70
Ξ	-63	-57
cutoff	800	525

Decay	8s	$\chi(2)$
$N(1710) \rightarrow N\pi\pi$	25	25
$N(1710) \rightarrow N\eta\pi$	0.58	0.32
$\Sigma(1770) \rightarrow N\bar{K}\pi$	4.7	4.5
$\Sigma(1770) \rightarrow \Sigma\pi\pi$	10	3.6
$\Xi(1860) \rightarrow \Sigma\bar{K}\pi$	0.57	0.40

Almost the same results

Difference comes from the SU(3) breaking of momenta at the vertex

27 and mass Lagrangians

$$\mathcal{L}^{27} = \frac{g^{27}}{2f} \left[4\bar{P}_{ijk}\epsilon^{lbk}\phi_l{}^i\phi_a{}^jB_b{}^a - \frac{4}{5}\bar{P}_{ijk}\epsilon^{lbk}\phi_l{}^a\phi_a{}^jB_b{}^i \right] + h.c.$$

$$\mathcal{L}^M = \frac{g^M}{2f} \bar{P}_{ijk}\epsilon^{lmk} \left(-\frac{1}{2f^2} \right) (2\phi M\phi + \phi\phi M + M\phi\phi)_l{}^iB_m{}^j + h.c.$$

**Fitting couplings to the N(1710) decay
-> large binding energy of 1 GeV : unrealistic**


Treat them as a small perturbation to the 8s.

$$g^{27} = g^M = g^{8s} = 1.88 , \quad b_{27} = -\frac{5}{4}(1-a) , \quad b_M = \frac{f^2}{m_\pi^2}(1-a)$$


$$\mathcal{L}^{int} = a\mathcal{L}^{8s} + b_{27,M}\mathcal{L}^{27,M}$$

Deviation from $a = 1$: weight of new terms

Results of 27 and mass Lagrangians



27
 $0.90 < a < 1.06$



M
 $0.76 < a < 1.06$

Contributions of these terms are considered as a theoretical uncertainty in the analysis.

Conclusion 1 : self-energy

We study the two-meson virtual cloud effect to the self-energy of baryon antidecuplet.



Two types of Lagrangians (8s, 8a) are important among several possible interaction Lagrangians.

$$\mathcal{L}^{8s} = \frac{g^{8s}}{2f} \bar{P}_{ijk} \epsilon^{lmk} \phi_l^a \phi_a^i B_m^j + h.c.$$

$$\mathcal{L}^{8a} = i \frac{g^{8a}}{4f^2} \bar{P}_{ijk} \epsilon^{lmk} \gamma^\mu (\partial_\mu \phi_l^a \phi_a^i - \phi_l^a \partial_\mu \phi_a^i) B_m^j + h.c.$$

Conclusion 1 : self-energy





- Two-meson cloud effects are always attractive, and contribute to the antidecuplet mass splitting, of the order of 20%.
- Antidecuplet members have relatively small decay widths to MMB channel.

A. Hosaka, T. H., F. J. Llanes-Estrada, E. Oset, J. R. Pelaez,
M. J. Vicente Vacas, hep-ph/0411311

Results of reaction : cross sections

Total cross section of



Conclusion 2 : reactions

We investigate the Θ production in (π^-, K^-) and (K^+, π^+) reactions, with the vertices obtained from the self-energy study.



The small cross section of the order of a few micro barn in (π^-, K^-) reaction may require some special mechanisms, such as interference of two amplitudes.



Self-energy

**Chiral symmetric Lagrangian,
Possible mixing with the other flavor
multiplets (8, 27, ...),**



Reaction

**Quantitative analysis (Form factor),
background cross section**