



# Meson production and baryon resonance

T. Sato (Osaka Univ.)

- Nucleon resonance study from meson production
- Pion photoproduction above 2GeV  
Non-perturbative to perturbative  
induced polarization  $p_y$

# Meson photoproduction and nucleon resonance





Thoma, Nucl-ex/0501024

Pion production →  $N^*$ , Delta resonances




$\Lambda^* + K$  production → missing  $N^*$

It may be more interesting to use open strangeness to study  $\Lambda^*$

Use  $q^2 Q$  system to investigate color  $3^*$ ,  $B=2/3$  system

# Importance of coupled channel effects

N-Delta quadrupole transition



Coupled channel study must be done to extract N\* paramters

K-matrix method Giessen group  
nucl-th/0505010

Cutkosky Model Pitt-FSU

Dynamical model DMT, Shizuoka-Osaka-ANL(SOA)


SOA: pi pi N three body unitarity

→ systematic data set

# Transition from Non-perturbative to perturbative?

$$\theta_{cm} = 90^\circ$$



Onset of scaling?



$$\vec{\gamma} + p \rightarrow \pi^0 + \vec{p}$$

$$Cx, Cz, py$$

### Induced polarization $p_y$



High spin state  
of baryon?

N			Delta	
	-	+	-	+
				2950 K15 **
		2700 K13 **	2750 I13 **	
2600 I 11	**			2420 H11 ****
2250G 9	****	2220 H 9 ****	2400G 9 **	2300 H 9 **
2190G 7	****		2200G 7 *	2390 F 7 *
				1950 F 7 ***

Lambda			Sigma	
	-	+	-	+
		2350 H 9 ***		
2100 G 7 ****		2020 F 7 *	2100 G 7 *	2030 F 7 ****

PDG

- Nucleon resonance study from meson production
- Pion photoproduction above 2GeV  
Non-perturbative to perturbative  
induced polarization  $p_y$