

PROPOSAL FOR EXPERIMENT AT RCNP

22 January 2004

TITLE: Continuing Investigation of the Compressional-mode Giant Resonances: Measurements in the Sn Isotopes as Tests for the Non-relativistic and Relativistic Calculations for the Nuclear Incompressibility

SPOKESPERSONS:

U. Garg, Physics Department, University of Notre Dame, Notre Dame, IN 46556, USA

e-mail: garg@nd.edu

FAX: 1.574.631.5952

M. Fujiwara, Research Center for Nuclear Physics, Osaka University, Suita, Osaka 567-0047, Japan

e-mail: fujiwara@rcnp.osaka-u.ac.jp

EXPERIMENTAL GROUPS:

University of Notre Dame, USA - Kyoto University, Japan - Research Center for Nuclear Physics, Japan - KVI, the Netherlands.

RUNNING TIME:

Total running time not including beam preparation 8 days

BEAM LINE:**BEAM REQUIREMENTS:**

Type of particle ^4He

Beam energy 400 MeV

Beam intensity 2 nA

Other requirements beam must be halo-free

highest stability over several days is required

BUDGET: Summary of budget expenses

Experimental expenses 2,000,000 yen

TITLE:

Continuing Investigation of the Compressional-mode Giant Resonances: Measurements in the Sn Isotopes as Tests for the Non-relativistic and Relativistic Calculations for the Nuclear Incompressibility

SPOKESPERSONS:

U. Garg, Physics Department, University of Notre Dame, Notre Dame, IN 46556, USA

M. Fujiwara, Research Center for Nuclear Physics, Osaka University, Suita, Osaka 567-0047, Japan

SUMMARY OF THE PROPOSAL

The proposed measurements aim at continuing detailed investigations on the compressional-mode giant resonances: the Giant Monopole Resonance and the Isoscalar Giant Dipole Resonance (ISGDR). Over the past few years, we have investigated the ISGDR at RCNP, using inelastic scattering of 400 MeV α particles at very small angles, including 0° . Our measurements, with the background-free spectra obtained with Grand Raiden, have resolved a longstanding problem and established that both the compressional-mode resonances (GMR and ISGDR) lead to the *same* value of nuclear incompressibility. However, there still remains a controversy in that the values of nuclear incompressibility obtained from relativistic and non-relativistic calculations based on the available GMR and ISGDR data are substantially different. We hope to help resolve this controversy by performing measurements on a series of Sn isotopes ($A=112-124$) to determine the centroids of the GMR strengths to about 100 keV, thereby providing constraints on the “symmetry” term in the expression of nuclear incompressibility.