

- [1] *E Aguayo, M Amman, F T Avignone, A S Barabash, P J Barton, J R Beene, F E Bertrand, M Boswell, V Brudanin, M Busch, Y.-D. Chan, C D Christofferson, J I Collar, D C Combs, R J Cooper, J A Detwiler, P J Doe, Y Efremenko, V Egorov, H Ejiri, S R Elliott, J Esterline, J E Fast, N Fields, P Finnerty, F M Fraenkle, A Galindo-Uribarri, V M Gehman, G K Giovanetti, M P Green, V E Guiseppe, K Gusey, A L Hallin, R Hazama, R Henning, E W Hoppe, M Horton, S Howard, M A Howe, R A Johnson, K J Keeter, M F Kidd, A Knecht, O Kochetov, S I Konovalov, R T Kouzes, B D LaFerriere, J Leon, L E Leviner, J C Loach, Q Looker, P N Luke, S MacMullin, M G Marino, R D Martin, J H Merriman, M L Miller, L Mizouni, M Nomachi, J L Orrell, N R Overman, G Perumpilly, D G Phillips, A W P Poon, D C Radford, K Rielage, R G H Robertson, M C Ronquest, A G Schubert, T Shima, M Shirchenko, K J Snavely, D Steele, J Strain, V Timkin, W Tornow, R L Varner, K Vetter, K Vorren, J F Wilkerson, E Yakushev, H Yaver, A R Young, C.-H. Yu, and V Yumatov.* 701:176–185, 2012, Characteristics of signals originating near the lithium-diffused N⁺ contact of high purity germanium p-type point contact detectors.
- [2] *J K Ahn.* *Few-Body Syst.*, 54(7-10):1043–1046, 2013, Photoproduction of Excited Hyperons Near 1,670 MeV/c².
- [3] *J K Ahn, H Akikawa, S Aoki, K Arai, S Y Bahk, K M Baik, B Bassalleck, J H Chung, M S Chung, D H Davis, T Fukuda, K Hoshino, A Ichikawa, M Ieiri, K Imai, K Itonaga, Y H Iwata, Y S Iwata, H Kanda, M Kaneko, T Kawai, M Kawasaki, C O Kim, J Y Kim, S H Kim, S J Kim, Y Kondo, T Kouketsu, H N Kyaw, Y L Lee, J W C McNabb, A A Min, M Mitsuhashi, K Miwa, K Nakazawa, Y Nagase, C Nagoshi, Y Nakanishi, H Noumi, S Ogawa, H Okabe, K Oyama, B D Park, H M Park, I G Park, J Parker, Y S Ra, J T Rhee, A Rusek, A Sawa, H Shibuya, K S Sim, P K Saha, D Seki, M Sekimoto, J S Song, H Takahashi, T Takahashi, F Takeuchi, H Tanaka, K Tanida, K T Tint, J Tojo, H Torii, S Torikai, D N Tovee, T Tsunemi, M Ukai, N Ushida, T Wint, K Yamamoto, N Yasuda, J T Yang, C J Yoon, C S Yoon, M Yosoi, T Yoshida, and L Zhu.* *Phys. Rev. C - Nucl. Phys.*, 88(1), 2013, Double- Λ hypernuclei observed in a hybrid emulsion experiment.
- [4] *S Ajimura, G Beer, H Bhang, M Bragadireanu, P Buehler, L Busso, M Cargnelli, S Choi, C Curceanu, S Enomoto, D Faso, H Fujioka, Y Fujiwara, T Fukuda, C Guaraldo, T Hashimoto, R S Hayano, T Hiraiwa, M Iio, M Iliescu, K Inoue, Y Ishiguro, T Ishikawa, S Ishimoto, T Ishiwatari, K Itahashi, M Iwai, M Iwasaki, S Kawasaki, P Kienle, H Kou, Y Ma, J Marton, Y Matsuda, Y Mizoi, O Morra, T Nagae, H Noumi, H Ohnishi, S Okada, H Outa, K Piscicchia, M P Lener, A R Vidal, Y Sada, A Sakaguchi, F Sakuma, M Sato, A Scordo, M Sekimoto, H Shi, D Sirghi, F Sirghi, K Suzuki, S Suzuki, T Suzuki, H Tatsuno, M Tokuda, D Tomono, A Toyoda, K Tsukada, O V Doce, E Widmann, T Yamazaki, H Yim, and J Zmeskal.* *Few-Body Syst.*, 54(7-10):1195–1199, 2013, A Search for Deeply Bound Kaonic Nuclear States at J-PARC.
- [5] *S Ajimura, G Beer, H Bhang, M Bragadireanu, P Buehler, L Busso, M Cargnelli, S Choi, C Curceanu, S Enomoto, D Faso, H Fujioka, Y Fujiwara, T Fukuda, C Guaraldo, T Hashimoto, R S Hayano, T Hiraiwa, M Iio, M Iliescu, K Inoue, Y Ishiguro, T Ishikawa, S Ishimoto, T Ishiwatari, K Itahashi, M Iwai, M Iwasaki, S Kawasaki, P Kienle, H Kou, Y Ma, J Marton, Y Matsuda, Y Mizoi, O Morra, T Nagae, H Noumi,*

H Ohnishi, S Okada, H Outa, K Piscicchia, M Poli Lener, A Romero Vidal, Y Sada, A Sakaguchi, F Sakuma, M Sato, A Scordo, M Sekimoto, H Shi, D Sirghi, F Sirghi, K Suzuki, S Suzuki, T Suzuki, H Tatsuno, M Tokuda, D Tomono, A Toyoda, K Tsukada, O Vazquez Doce, E Widmann, T Yamazaki, H Yim, and J Zmeskal. Nucl. Phys. A, 914:315–320, 2013, A search for deeply-bound kaonic nuclear state at the J-PARC E15 experiment.

- [6] *H Al Falou, R Kanungo, C Andreoiu, D S Cross, B Davids, M Djongolov, A T Gallant, N Galinski, D Howell, R Kshetri, D Niamir, J N Orce, A C Shotter, S Sjue, I Tanihata, I J Thompson, S Triambak, M Uchida, P Walden, and R B Wiringa. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys., 721(4-5):224–228, 2013, Inelastic scattering of ^9Li and excitation mechanism of its first excited state.*
- [7] *S Almaraz-Calderon, W P Tan, A Aprahamian, M Beard, G P A Berg, B Bucher, M Couder, J Görres, S O’Brien, D Patel, A Roberts, K Sault, M Wiescher, C R Brune, T N Massey, K Fujita, K Hatanaka, D Ishiwaka, H Matsubara, H Okamura, H J Ong, Y Sakemi, Y Shimizu, T Suzuki, Y Tameshige, A Tamii, J Zenihiro, T Kubo, Y Namiki, Y Ohkuma, Y Shimbara, S Suzuki, R Watanabe, R Yamada, T Adachi, Y Fujita, H Fujita, M Dozono, and T Wakasa. Phys. Rev. C - Nucl. Phys., 88(5), 2013, Erratum: Level structure of ^{30}S and its importance in the $^{26}\text{Si}(\alpha, p)^{29}\text{P}$ and $^{29}\text{P}(p, \gamma)^{30}\text{S}$ reaction rates (Physical Review C - Nuclear Physics (2012) 86 (065805)).*
- [8] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol., 88(11), 2013, Search for charginos nearly mass degenerate with the lightest neutralino based on a disappearing-track signature in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector.*
- [9] *ATLAS Collaboration. Phys. Rev. Lett., 111(23), 2013, Measurement of top quark polarization in top-antitop events from proton-proton collisions at $\sqrt{s}=7$ TeV using the ATLAS detector.*
- [10] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol., 88(11), 2013, Search for long-lived stopped R-hadrons decaying out of time with pp collisions using the ATLAS detector.*
- [11] *ATLAS Collaboration. Eur. Phys. J. C, 73(12):1–31, 2013, Measurement of jet shapes in top-quark pair events at $\sqrt{s} = 7$ TeV using the ATLAS detector.*
- [12] *ATLAS Collaboration. J. High Energy Phys., 2013(10):1–37, 2013, Measurement of the differential cross-section of B^+ meson production in pp collisions at $\sqrt{s} = 7$ TeV at ATLAS.*
- [13] *ATLAS Collaboration. J. High Energy Phys., 2013(11), 2013, Measurement of the distributions of event-by-event flow harmonics in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector at the LHC.*
- [14] *ATLAS Collaboration. Nucl. Phys. B, 875(3):483–535, 2013, Dynamics of isolated-photon plus jet production in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector.*
- [15] *ATLAS Collaboration. Phys. Rev. Lett., 111(15), 2013, Measurement of the azimuthal angle dependence of inclusive jet yields in Pb+Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV with the ATLAS detector.*
- [16] *ATLAS Collaboration. J. High Energy Phys., 2013(10), 2013, Search for direct third-generation squark pair production in final states with missing transverse momentum and two b-jets in $\sqrt{s} = 8\text{TeV}$ pp collisions with the ATLAS detector.*

- [17] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 88(7), 2013, Search for microscopic black holes in a like-sign dimuon final state using large track multiplicity with the ATLAS detector.
- [18] *ATLAS Collaboration. J. High Energy Phys.*, 2013(10), 2013, Search for new phenomena in final states with large jet multiplicities and missing transverse momentum at $\sqrt{s} = 8$ TeV proton-proton collisions using the ATLAS experiment.
- [19] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 725(4-5):223–242, 2013, Measurement of the high-mass Drell-Yan differential cross-section in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector.
- [20] *ATLAS Collaboration. J. High Energy Phys.*, 2013(9), 2013, Performance of jet sub-structure techniques for large-R jets in proton-proton collisions at $\sqrt{s}=7$ TeV using the ATLAS detector.
- [21] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 726(1-3):88–119, 2013, Measurements of Higgs boson production and couplings in diboson final states with the ATLAS detector at the LHC.
- [22] *ATLAS Collaboration. J. High Energy Phys.*, 2013(2):1–46, 2013, Search for the neutral Higgs bosons of the minimal supersymmetric standard model in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [23] *ATLAS Collaboration. New J. Phys.*, 15, 2013, Search for excited electrons and muons in $\sqrt{s}=8$ TeV proton-proton collisions with the ATLAS detector.
- [24] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 726(1-3):120–144, 2013, Evidence for the spin-0 nature of the Higgs boson using ATLAS data.
- [25] *ATLAS Collaboration. J. High Energy Phys.*, 2013(7), 2013, Measurement of the production cross section of jets in association with a Z boson in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector.
- [26] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 725(1-3):60–78, 2013, Measurement with the ATLAS detector of multi-particle azimuthal correlations in p + Pb collisions at $\sqrt{s_{NN}}= 5.02$ TeV.
- [27] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 88(3), 2013, Measurement of charged-particle event shape variables in inclusive $\sqrt{s}=7$ TeV proton-proton interactions with the ATLAS detector.
- [28] *ATLAS Collaboration. Eur. Phys. J. C*, 73(8):1–56, 2013, Measurement of the inclusive jet cross-section in pp collisions at $\sqrt{s} = 2.76$ TeV and comparison to the inclusive jet cross-section at $\sqrt{s} = 7$ TeV using the ATLAS detector.
- [29] *ATLAS Collaboration. Eur. Phys. J. C*, 73(8):1–39, 2013, Improved luminosity determination in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector at the LHC.
- [30] *ATLAS Collaboration. J. High Energy Phys.*, 2013(6), 2013, Measurement of the cross-section for W boson production in association with b-jets in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector.
- [31] *ATLAS Collaboration. J. High Energy Phys.*, 2013(6), 2013, Search for third generation scalar leptoquarks in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector.

- [32] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 88(1), 2013, Search for tt resonances in the lepton plus jets final state with ATLAS using 4.7 fb-1 of pp collisions at $\sqrt{s}=7$ TeV.
- [33] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 88(1), 2013, Search for nonpointing photons in the diphoton and ETmiss final state in $\sqrt{s}=7$ TeV proton-proton collisions using the ATLAS detector.
- [34] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(11), 2013, Search for resonant diboson production in the WW/WZ ν jj decay channels with the ATLAS detector at $\sqrt{s}=7$ TeV.
- [35] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(11), 2013, Measurements of $W\gamma$ and $Z\gamma$ production in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector at the LHC.
- [36] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(11), 2013, Measurement of $W+W^-$ production in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector and limits on anomalous WWZ and $WW\gamma$ couplings.
- [37] *ATLAS Collaboration. Eur. Phys. J. C*, 73(6):1–20, 2013, Search for a light charged Higgs boson in the decay channel $H^\pm \rightarrow cs$ in tt events using pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [38] *ATLAS Collaboration. Phys. Rev. Lett.*, 110(18), 2013, Observation of Associated Near-Side and Away-Side Long-Range Correlations in $\sqrt{s_{NN}}=5.02$ TeV Proton-Lead Collisions with the ATLAS Detector.
- [39] *ATLAS Collaboration. Eur. Phys. J. C*, 73(5):1–30, 2013, Measurement of k_T splitting scales in $W \rightarrow \nu$ events at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [40] *ATLAS Collaboration. Eur. Phys. J. C*, 73(3):1–27, 2013, Jet energy resolution in proton-proton collisions at $\sqrt{s} = 7$ TeV recorded in 2010 with the ATLAS detector.
- [41] *ATLAS Collaboration. Eur. Phys. J. C*, 73(3):1–118, 2013, Jet energy measurement with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV.
- [42] *ATLAS Collaboration. Eur. Phys. J. C*, 73(3):1–33, 2013, Multi-channel search for squarks and gluinos in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector at the LHC.
- [43] *ATLAS Collaboration. Eur. Phys. J. C*, 73(3):1–34, 2013, Single hadron response measurement and calorimeter jet energy scale uncertainty with the ATLAS detector at the LHC.
- [44] *ATLAS Collaboration. Eur. Phys. J. C*, 73(3):1–18, 2013, Measurement of the tt production cross section in the tau + jets channel using the ATLAS detector.
- [45] *ATLAS Collaboration. J. High Energy Phys.*, 2013(4), 2013, Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the ATLAS detector.
- [46] *ATLAS Collaboration. Eur. Phys. J. C*, 73(2):1–30, 2013, Measurement of the flavour composition of dijet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [47] *ATLAS Collaboration. J. High Energy Phys.*, 2013(3), 2013, Search for charged Higgs bosons through the violation of lepton universality in tt events using pp collision data at $\sqrt{S}=7$ TeV with the ATLAS experiment.

- [48] *ATLAS Collaboration. J. High Energy Phys.*, 2013(3), 2013, Measurement of ZZ production in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous ZZZ and ZZ γ couplings with the ATLAS detector.
- [49] *ATLAS Collaboration. New J. Phys.*, 15, 2013, Search for extra dimensions in diphoton events from proton-proton collisions at $\sqrt{s} = 7$ TeV in the ATLAS detector at the LHC.
- [50] *ATLAS Collaboration. New J. Phys.*, 15, 2013, Search for WH production with a light Higgs boson decaying to prompt electron-jets in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [51] *ATLAS Collaboration. Eur. Phys. J. C*, 73(1):1–20, 2013, Search for pair-produced massive coloured scalars in four-jet final states with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV.
- [52] *ATLAS Collaboration. Eur. Phys. J. C*, 73(1):1–28, 2013, Measurements of top quark pair relative differential cross-sections with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV.
- [53] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(5), 2013, Measurement of upsilon production in 7 TeV pp collisions at ATLAS.
- [54] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(5), 2013, Search for new phenomena in events with three charged leptons at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [55] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 719(4-5):299–317, 2013, A search for prompt lepton-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [56] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 719(4-5):261–279, 2013, Search for supersymmetry in events with photons, bottom quarks, and missing transverse momentum in proton-proton collisions at a centre-of-mass energy of 7 TeV with the ATLAS detector.
- [57] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 719(4-5):220–241, 2013, Measurement of the jet radius and transverse momentum dependence of inclusive jet suppression in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector.
- [58] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 719(4-5):242–260, 2013, A search for high-mass resonances decaying to $\tau + \tau^-$ in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [59] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 719(4-5):280–298, 2013, Search for long-lived, heavy particles in final states with a muon and multi-track displaced vertex in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [60] *ATLAS Collaboration. J. High Energy Phys.*, 2013(1), 2013, Search for direct chargino production in anomaly-mediated supersymmetry breaking models based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.
- [61] *ATLAS Collaboration. J. High Energy Phys.*, 2013(1), 2013, Search for resonances decaying into top-quark pairs using fully hadronic decays in pp collisions with ATLAS at $\sqrt{s} = 7$ TeV.
- [62] *ATLAS Collaboration. J. High Energy Phys.*, 2013(1), 2013, Measurement of isolated-photon pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector.

- [63] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(3), 2013, Measurement of the Λ_b^0 lifetime and mass in the ATLAS experiment.
- [64] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 718(4-5):1284–1302, 2013, Search for pair production of heavy top-like quarks decaying to a high-pT W boson and a b quark in the lepton plus jets final state at $s=7\text{TeV}$ with the ATLAS detector.
- [65] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(1), 2013, Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb⁻¹ of $s=7\text{ TeV}$ proton-proton collision data.
- [66] *ATLAS Collaboration. Phys. Rev. Lett.*, 110(2), 2013, Measurement of Z boson production in Pb-Pb collisions at $\sqrt{s_{NN}}=2.76\text{ TeV}$ with the ATLAS detector.
- [67] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 718(3):841–859, 2013, Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $s=7\text{ TeV}$ pp collisions with the ATLAS detector.
- [68] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 718(3):860–878, 2013, Search for new phenomena in the WW $l\nu l'\nu'$ final state in pp collisions at $s=7\text{TeV}$ with the ATLAS detector.
- [69] *ATLAS Collaboration. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 718(3):879–901, 2013, Search for direct slepton and gaugino production in final states with two leptons and missing transverse momentum with the ATLAS detector in pp collisions at $s=7\text{ TeV}$.
- [70] *ATLAS Collaboration. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(1), 2013, Search for contact interactions and large extra dimensions in dilepton events from pp collisions at $s=7\text{ TeV}$ with the ATLAS detector.
- [71] *ATLAS Collaboration. Phys. Rev. Lett.*, 110(1), 2013, Search for dark matter candidates and large extra dimensions in events with a photon and missing transverse momentum in pp collision data at $s=7\text{ TeV}$ with the ATLAS detector.
- [72] *ATLAS Collaboration. J. High Energy Phys.*, 2013(11), 2013, Measurement of the top quark charge in pp collisions at $s = 7\text{ TeV}$ with the ATLAS detector.
- [73] *L Audirac, A Obertelli, P Doornenbal, D Mancusi, S Takeuchi, N Aoi, H Baba, S Boissinot, A Boudard, A Corsi, A Gillibert, T Isobe, A Jungclaus, V Lapoux, J Lee, S Leray, K Matsui, M Matsushita, T Motobayashi, D Nishimura, S Ota, E C Pollacco, G Potel, H Sakurai, C Santamaria, Y Shiga, D Sohler, D Steppenbeck, R Taniuchi, and H Wang. Phys. Rev. C - Nucl. Phys.*, 88(4), 2013, Evaporation-cost dependence in heavy-ion fragmentation.
- [74] *T Aumann, K Blaum, Y Blumenfeld, A Bonaccorso, M J G Borge, L Brink, A Sharpy, J Dilling, G D Dracoulis, C Fahlander, C Forssén, H O U Fynbo, A Gade, P T Greenlees, F Gulminelli, K Heyde, A Heinz, M Hjorth-Jensen, R V F Janssens, K L Jones, B Jonson, R Kanungo, E Khan, K Langanke, T Nakamura, A Navin, W Nazarewicz, T Nilsson, G Nyman, W Nörtershäuser, T Otsuka, M Pfützner, K Rüsager, J Rotureau, E Ryberg, H Schatz, N Severijns, H Simon, O Sorlin, D Sääf, I Tanihata, R Thies, P Van Duppen, and M Zhukov. T152*, 2013, Nobel symposium 152: Physics with radioactive beams.

- [75] *H Bhang, S Ajimura, K Aoki, A Banu, T Fukuda, O Hashimoto, J I Hwang, S Kameoka, B H Kang, E Kim, J H Kim, M Kim, T Maruta, Y Miura, Y Miyake, T Nagae, M Nakamura, S N Nakamura, H Noumi, S Okada, Y Okayasu, H Outa, H Park, P K Saha, Y Sato, M Sekimoto, T Takahashi, H Tamura, K Tanida, A Toyoda, K Tshoo, K Tsukada, T Watanabe, and H J Yim. Few-Body Syst.*, 54(1-4):103–110, 2013, Three-Body ANN nNN Nonmesonic Weak Decay Process of Λ Hypernuclei.
- [76] *H Bhang, S Ajimura, K Aoki, A Banu, T Fukuda, O Hashimoto, J I Hwang, S Kameoka, B H Kang, E Kim, J H Kim, M Kim, T Maruta, Y Miura, Y Miyake, T Nagae, M Nakamura, S N Nakamura, H Noumi, S Okada, Y Okayasu, H Outa, H Park, P K Saha, Y Sato, M Sekimoto, T Takahashi, H Tamura, K Tanida, A Toyoda, K Tshoo, K Tsukada, T Watanabe, and H J Yim. Few-Body Syst.*, 54(7-10):1239–1243, 2013, The Strong Three-body Weak Interaction Contribution in the Nonmesonic Weak Decay of p-shell Λ Hypernuclei.
- [77] *M Bowry, Zs. Podolyák, S Pietri, J Kurcewicz, M Bunce, P H Regan, F Farinon, H Geissel, C Nociforo, A Prochazka, H Weick, N Al-Dahan, N Alkhomashi, P R P Allegro, J Benlliure, G Benzoni, P Boutachkov, A M Bruce, A M Denis Bacelar, G F Farrelly, J Gerl, M Górska, A Gottardo, J Grbosz, N Gregor, R Janik, R Knöbel, I Kojouharov, T Kubo, N Kurz, Y A Litvinov, E Merchan, I Mukha, F Naqvi, B Pfeiffer, M Pfützner, W Plaß, M Pomorski, B Riese, M V Ricciardi, K.-H. Schmidt, H Schaffner, C Scheidenberger, E C Simpson, B Sitar, P Spiller, J Stadlmann, P Strmen, B Sun, I Tanihata, S Terashima, J J Valiente Dobón, J S Winfield, H.-J. Wollersheim, and P J Woods. Phys. Rev. C - Nucl. Phys.*, 88(2), 2013, Population of high-spin isomeric states following fragmentation of ^{238}U .
- [78] *W.-C. Chang. Few-Body Syst.*, 54(1-4):171–176, 2013, Recent Results from LEPS/SPring-8 Experiment.
- [79] *S Cook, R D’Arcy, M Fukuda, K Hatanaka, Y Hino, Y Kuno, M Lancaster, Y Mori, T H Nam, T Ogitsu, H Sakamoto, A Sato, N M Truong, A Yamamoto, M Yoshida, and M Wing. 408(1)*, 2013, First measurements of muon production rate using a novel pion capture system at MuSIC.
- [80] *P Doornenbal, H Scheit, S Takeuchi, N Aoi, K Li, M Matsushita, D Steppenbeck, H Wang, H Baba, H Crawford, C R Hoffman, R Hughes, E Ideguchi, N Kobayashi, Y Kondo, J Lee, S Michimasa, T Motobayashi, H Sakurai, M Takechi, Y Togano, R Winkler, and K Yoneda. Phys. Rev. Lett.*, 111(21), 2013, In-beam γ -ray spectroscopy of $\text{Mg}^{34,36,38}$: Merging the $N=20$ and $N=28$ shell quenching.
- [81] *H Ejiri, I H Hashim, Y Hino, Y Kuno, Y Matsumoto, K Ninomiya, H Sakamoto, A Sato, T Shima, A Shinohara, K Takahisa, and N H Tran. J. Phys. Soc. Japan*, 82(4), 2013, Nuclear γ rays from stopped muon capture reactions for nuclear isotope detection.
- [82] *X Fang, B Bucher, S Almaraz-Calderon, A Alongi, A D Ayangeakaa, A Best, G P A Berg, C Cahillane, E Dahlstrom, R J Deboer, M Freer, H Fujita, Y Fujita, J Görres, K Hatanaka, A Howard, T Itoh, T Kadoya, T Kawabata, J J Kolata, Q Li, Y J Li, B Liu, A Long, Y.-W. Lui, S Lyons, Y Matsuda, K Miki, N Paul, A Roberts, M K Smith, R Talwar, A Tamii, W P Tan, X D Tang, M Wiescher, and N Yokota. 420(1)*, 2013, Experimental investigation of the $^{12}\text{C}+^{12}\text{C}$ fusion at very low energies by direct and indirect methods.
- [83] *H. Fujita, Y. Fujita, T. Adachi, H. Akimune, N. T. Botha, K. Hatanaka, H. Matsubara, K. Nakanishi, R. Neveling, A. Okamoto, Y. Sakemi, T. Shima, Y. Shimizu, F. D. Smit,*

T. Suzuki, A. Tamii, and M. Yosoi. Phys. Rev. C, 88(5):054329, November 2013, Isospin mixing of the isobaric analog state studied in a high-resolution $^{56}\text{Fe}(^3\text{He},t)^{56}\text{Co}$ reaction.

- [84] *H Fujita, Y Fujita, T Adachi, H Akimune, N T Botha, K Hatanaka, H Matsubara, K Nakanishi, R Neveling, A Okamoto, Y Sakemi, T Shima, Y Shimizu, F D Smit, T Suzuki, A Tamii, and M Yosoi. Phys. Rev. C - Nucl. Phys.*, 88(5), 2013, Isospin mixing of the isobaric analog state studied in a high-resolution $^{56}\text{Fe}(^3\text{He},t)^{56}\text{Co}$ reaction.
- [85] *Y Fujita, T Adachi, H Fujita, A Algora, B Blank, M Csatlós, J M Deaven, E Estevez-Aguado, E Ganiolu, C J Guess, J Gulyás, K Hatanaka, K Hirota, M Honma, D Ishikawa, A Krasznahorkay, H Matsubara, R Meharchand, F Molina, H Okamura, H J Ong, T Otsuka, G Perdikakis, B Rubio, C Scholl, Y Shimbara, E J Stephenson, G Susoy, T Suzuki, A Tamii, J H Thies, R G T Zegers, and J Zenihiro. Phys. Rev. C - Nucl. Phys.*, 88(1), 2013, High-resolution study of $T_z=+2 \rightarrow +1$ Gamow-Teller transitions in the $^{44}\text{Ca}(^3\text{He},t)^{44}\text{Sc}$ reaction.
- [86] *K Fushimi, S Nakayama, R Orito, R Sugawara, Y Awatani, H Ejiri, T Shima, R Hazama, K Inoue, H Ikeda, and A Kozlov. 469(1)*, 2013, PICO-LON dark matter search.
- [87] *E Ganiolu, H Fujita, Y Fujita, T Adachi, A Algora, M Csatlós, J M Deaven, E Estevez-Aguado, C J Guess, J Gulyás, K Hatanaka, K Hirota, M Honma, D Ishikawa, A Krasznahorkay, H Matsubara, R Meharchand, F Molina, H Okamura, H J Ong, T Otsuka, G Perdikakis, B Rubio, C Scholl, Y Shimbara, G Susoy, T Suzuki, A Tamii, J H Thies, R G T Zegers, and J Zenihiro. Phys. Rev. C - Nucl. Phys.*, 87(1), 2013, High-resolution study of Gamow-Teller transitions in the $^{47}\text{Ti}(^3\text{He},t)^{47}\text{V}$ reaction.
- [88] *K Hosomi, S Ajimura, K Aoki, M Dairaku, Y Y Fu, H Fujioka, T Fukuda, K Futatsukawa, W Imoto, Y Kakiguchi, M Kawai, S Kinoshita, T Koike, Y Ma, N Maruyamai, M Mimori, S Minami, Y Miura, K Miwa, Y Miyagi, T Nagae, D Nakajima, H Noumi, K Shirotori, T Suzuki, T Takahashi, T N Takahashi, H Tamura, K Tanida, N Terada, A Toyoda, K Tsukada, M Ukai, and S H Zhou. Nucl. Phys. A*, 914:184–188, 2013, Gamma-ray spectroscopy of $^{12}\Lambda\text{C}$ via the (π^+, K^+) reaction.
- [89] *K Hosomi, S Ajimura, K Aoki, M Dairaku, Y Y Fu, H Fujioka, T Fukuda, K Futatsukawa, W Imoto, Y Kakiguchi, M Kawai, S Kinoshita, T Koike, Y Ma, N Maruyamai, M Mimori, S Minami, Y Miura, K Miwa, Y Miyagi, T Nagae, D Nakajima, H Noumi, K Shirotori, T Suzuki, T Takahashi, T N Takahashi, H Tamura, K Tanida, N Terada, A Toyoda, K Tsukada, M Ukai, and S H Zhou. Nucl. Phys. A*, 914:184–188, 2013, Gamma-ray spectroscopy of $^{12}\Lambda\text{C}$ via the (π^+, K^+) reaction.
- [90] *J W Hwang, Y Satou, T Nakamura, Y Kondo, N Matsui, Y Hashimoto, T Nakabayashi, T Okumura, M Shinohara, N Fukuda, T Sugimoto, H Otsu, Y Togano, T Motobayashi, H Sakurai, Y Yanagisawa, N Aoi, S Takeuchi, T Gomi, M Ishihara, S Kawai, H J Ong, T K Onishi, S Shimoura, M Tamaki, T Kobayashi, Y Matsuda, N Endo, and M Kitayama. Few-Body Syst.*, 54(7-10):1469–1472, 2013, Discovery of the First 2- State in ^{16}C via Neutron Knockout Reaction.
- [91] *S Hwang, J K Ahn, and K Hicks. Few-Body Syst.*, 54(1-4):265–269, 2013, $K^*0\Sigma^+$ Photoproduction from Threshold to 3.0 GeV.
- [92] *S Hwang, K Hicks, J K Ahn, and T Nakano. Few-Body Syst.*, 54(7-10):1037–1041, 2013, $K^*0\Sigma^+$ Photoproduction with Evidence for the Kappa Meson Exchange at SPring 8/LEPS.

- [93] Y Ichikawa, H C Bhang, S Bufalino, H Ekawa, P Evtoukhovitch, A Feliciello, H Fujioka, S Hasegawa, S Hayakawa, R Honda, K Hosomi, K Imai, S Ishimoto, C W Joo, S Kanatsuki, R Kiuchi, T Koike, H Kumawat, Y Matsumoto, K Miwa, M Moritsu, T Nagae, M Naruki, M Niiyama, Y Nozawa, R Ota, A Sakaguchi, H Sako, V Samoilov, S Sato, K Shirotori, H Sugimura, S Suzuki, T Takahashi, T N Takahashi, H Tamura, T Tanaka, K Tanida, A O Tokiyasu, Z Tsamalaidze, B J Roy, M Ukai, T O Yamamoto, and S B Yang. *Few-Body Syst.*, 54(7-10):1191–1194, 2013, J-PARC E27 Experiment to Search for a Nuclear Kaon Bound State K-pp.
- [94] Y Ichikawa, H Ueno, Y Ishii, T Furukawa, A Yoshimi, D Kameda, H Watanabe, N Aoi, K Asahi, D L Balabanski, R Chevrier, J M Daugas, N Fukuda, G Georgiev, H Hayashi, H Iijima, N Inabe, T Inoue, M Ishihara, T Kubo, T Nanao, T Ohnishi, K Suzuki, M Tsuchiya, H Takeda, and M M Rajabali. *Hyperfine Interact.*, 220(1-3):47–51, 2013, Novel method for the production of spin-aligned RI beams in projectile fragmentation reaction with the dispersion matching technique.
- [95] Y Ichikawa, H Ueno, Y Ishii, T Furukawa, A Yoshimi, D Kameda, H Watanabe, N Aoi, K Asahi, D L Balabanski, R Chevrier, J M Daugas, N Fukuda, G Georgiev, H Hayashi, H Iijima, N Inabe, T Inoue, M Ishihara, T Kubo, T Nanao, T Ohnishi, K Suzuki, M Tsuchiya, H Takeda, and M M Rajabali. *Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms*, 317(PART B):769–773, 2013, Spin-aligned RI beams via two-step fragmentation reactions.
- [96] M Ieiri, M Minakawa, K Agari, E Hirose, Y Katoh, R Muto, M Naruki, H Noumi, Y Sato, S Sawada, Y Shirakabe, Y Suzuki, H Takahashi, M Takasaki, K H Tanaka, A Toyoda, H Watanabe, and Y Yamanoi. *Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms*, 317(PART B):338–341, 2013, Electrostatic separators in the hadron experimental facility at J-PARC.
- [97] M Itoh, H Akimune, M Fujiwara, U Garg, T Kawabata, K Kawase, T Murakami, K Nakanishi, Y Nakatsugawa, H Sakaguchi, S Terashima, M Uchida, Y Yasuda, M Yosoi, and J Zenihiro. 436(1), 2013, Nature of 10 MeV state in ^{12}C .
- [98] M Itoh, S Kishi, H Sakaguchi, H Akimune, M Fujiwara, U Garg, K Hara, H Hashimoto, J Hoffman, T Kawabata, K Kawase, T Murakami, K Nakanishi, B K Nayak, S Terashima, M Uchida, Y Yasuda, and M Yosoi. *Phys. Rev. C - Nucl. Phys.*, 88(6), 2013, Isoscalar giant resonance strengths in ^{32}S and possible excitation of superdeformed and $^{28}\text{Si} + \alpha$ cluster bandheads.
- [99] R Itoh, Y Oga, S Noguchi, H Igarashi, and H Ueda. *Phys. C Supercond. its Appl.*, 484:300–304, 2013, Numerical simulation of screening current distribution in HTS tape of high field magnet.
- [100] J Kamiya, N Ogiwara, M Kinsho, K Kamakura, N Hamatani, K Hatanaka, M Fukuda, and K Takahisa. *J. Vac. Soc. Japan*, 56(5):167–171, 2013, Titanium alloy as low activation vacuum material.
- [101] T Kawabata, T Adachi, M Fujiwara, K Hatanaka, Y Ishiguro, M Itoh, Y Maeda, H Matsu-Subara, H Miyasako, Y Nozawa, T Saito, S Sakaguchi, Y Sasamoto, Y Shimizu, T Takahashi, A Tamii, S Terashima, H Tokieda, N Tomida, T Uesaka, M Uchida, Y Yasuda, N Yokota, H P Yoshida, and J Zenihiro. *Few-Body Syst.*, 436(1-4):93–97, 2013, Search for alpha inelastic condensed state in ^{24}Mg .
- [102] H Kawamura, H Arikawa, S Ezure, K Harada, T Hayamizu, T Inoue, T Ishikawa, M Itoh, T Kato, T Sato, T Aoki, T Furukawa, A Hatakeyama, K Hatanaka, K Imai, T Murakami,

H S Nataraj, Y Shimizu, T Wakasa, H P Yoshida, and Y Sakemi. Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms, 317(PART B):582–585, 2013, Laser-cooled radioactive francium factory at CYRIC.

- [103] *T Kishimoto, F Khanam, T Hayakawa, S Ajimura, T Itabashi, K Matsuoka, S Minami, Y Mitoma, A Sakaguchi, Y Shimizu, K Terai, T Sato, H Noumi, M Sekimoto, H Takahashi, T Fukuda, W Imoto, and Y Mizoi. Prog. Theor. Exp. Phys.*, 2013(4), 2013, Search for a bound kaon and pion state.
- [104] *N Kobayashi, T Nakamura, Y Kondo, N Aoi, H Baba, S Deguchi, N Fukuda, G S Lee, H S Lee, N Inabe, M Ishihara, Y Kawada, R Kanungo, T Kubo, M A Famiano, M Matsushita, T Motobayashi, T Ohnishi, N A Orr, H Otsu, R Barthelemy, H Sakurai, S Kim, T Sako, T Sumikama, Y Satou, K Takahashi, H Takeda, M Takechi, S Takeuchi, K N Tanaka, N Tanaka, R Tanaka, Y Togano, and K Yoneda. Few-Body Syst.*, 54(7-10):1441–1444, 2013, Breakup Reactions of Drip-Line Nuclei Near $N = 20$, 28.
- [105] *T Kubota, T Kishida, O Jinnouchi, Y Ikegami, Y Unno, S Terada, S Mitsui, A Tamii, T Aoi, K Hanagaki, K Hara, N Kimura, R Takashima, Y Takubo, J Tojo, K Nagai, I Nakano, and K Yorita. Nucl. Instruments Methods Phys. Res. Sect. A Accel. Spectrometers, Detect. Assoc. Equip.*, 731:247–251, 2013, Beam test of novel n-in-p strip sensors for high radiation environment.
- [106] *S Kurashima, S Okumura, N Miyawaki, H Kashiwagi, T Satoh, T Kamiya, M Fukuda, and W Yokota. Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms*, 306:40–43, 2013, Short-time change of heavy-ion microbeams with different mass to charge ratios by scaling method for the JAEA AVF cyclotron.
- [107] *R Leguillon, H Nishibata, Y Ito, C M Petrache, A Odahara, T Shimoda, N Hamatani, K Tajiri, J Takatsu, R Yokoyama, E Ideguchi, H Watanabe, Y Wakabayashi, K Yoshinaga, T Suzuki, S Nishimura, D Beaumel, G Lehaut, D Guinet, P Desesquelles, D Curien, A Astier, T Konstantinopoulos, and T Zerrouki. Phys. Rev. C - Nucl. Phys.*, 88(4), 2013, Spectroscopy of ^{135}La .
- [108] *W Liu, K Hatanaka, S Zhu, and N Aoi. 1533:1*, 2013, Preface: 8th China-Japan joint nuclear physics symposium.
- [109] *Y Maeda, T Saito, H Miyasako, T Uesaka, S Ota, S Kawase, T Kikuchi, H Tokieda, T Kawabata, K Yako, T Wakasa, S Sakaguchi, R Chen, H Sakaguchi, T Shima, T Suzuki, and A Tamii. Few-Body Syst.*, 54(1-4):291–294, 2013, Measurements of the Cross Sections and $A\gamma$ for $D(p, n)$ Inclusive Breakup Reaction at 170 MeV.
- [110] *Y Maeda, T Saito, H Miyasako, T Uesaka, S Ota, S Kawase, T Kikuchi, H Tokieda, T Kawabata, K Yako, T Wakasa, S Sakaguchi, R Chen, H Sakaguchi, T Shima, T Suzuki, and A Tamii. Few-Body Syst.*, 54(7-10):1311–1314, 2013, Study of the Three-Nucleon Force Effects in the $2H(p, n)$ Breakup Reaction at 170 MeV.
- [111] *V Mares, C Pioch, W Ruhm, H Iwase, Y Iwamoto, M Hagiwara, D Satoh, H Yashima, T Itoga, T Sato, Y Nakane, H Nakashima, Y Sakamoto, T Matsumoto, A Masuda, H Harano, J Nishiyama, C Theis, E Feldbaumer, L Jaegerhofer, A Tamii, K Hatanaka, and T Nakamura. IEEE Trans. Nucl. Sci.*, 60(1):299–304, 2013, Neutron dosimetry in quasi-monoenergetic fields of 244 and 387 MeV.
- [112] *H Matsubara, M Takaki, T Uesaka, S Shimoura, N Aoi, M Dozono, T Fujii, K Hatanaka, T Hashimoto, T Kawabata, S Kawase, K Kisamori, Y Kikuch, Y Kubota, C S Lee, H C Lee, Y Maeda, S Michimasa, K Miki, H Miya, S Noji, S Ota, S Sakaguchi, Y Sasamoto,*

T Suzuki, L T Tang, K Takahisa, H Tokieda, A Tamii, K Yako, Y Yasuda, N Yokota, R Yokoyama, and J Zenihiro. Few-Body Syst., 54(7-10):1433–1436, 2013, Spectroscopic Measurement in ^9He and ^{12}Be .

- [113] *K Matsuta, Y Masuda, K Hatanaka, S C Jeong, S Kawasaki, R Matsumiya, M Mihara, Y Watanabe, D Nishimura, Y Morita, K Asahi, T Adachi, J Martin, A Konaka, A Miller, C Bidinosti, T Dawson, L Lee, C Davis, D Ramsay, W van Oers, E Korkmaz, and L Buchman. Hyperfine Interact., 220(1-3):89–93, 2013, Creation of polarized ultracold neutrons and observation of Ramsey resonance for electric dipole moment measurement.*
- [114] *K Matsuta, T Minamisono, M Mihara, M Fukuda, S Zhu, Y Masuda, K Hatanaka, D Yuan, Y Zheng, Y Zuo, P Fang, D Zhou, T Ohtsubo, T Izumikawa, S Momota, D Nishimura, R Matsumiya, A Kitagawa, S Sato, M Kanazawa, M Torikoshi, T Nagatomo, Y Nojiri, T Suzuki, X Zhang, J R Alonso, T J M Symons, S Kawasaki, S C Jeong, Y Watanabe, and K Minamisono. Hyperfine Interact., 220(1-3):21–28, 2013, Nuclear moments as a probe of electronic structure in material, exotic nuclear structure and fundamental symmetry.*
- [115] *S Michimasa, M Takaki, M Dozono, S Go, H Baba, E Ideguchi, K Kisamori, H Matsubara, H Miya, S Ota, H Sakai, S Shimoura, A Stolz, T L Tang, H Tokieda, T Uesaka, and R G T Zegers. Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms, 317(PART B):710–713, 2013, Development of CVD diamond detector for time-of-flight measurements.*
- [116] *K Miki, A Tamii, N Aoi, T Fukui, T Hashimoto, K Hatanaka, T Ito, T Kawabata, H Matsubara, K Ogata, H J Ong, H Sakaguchi, S Sakaguchi, T Suzuki, J Tanaka, I Tanihata, T Uesaka, and T Yamamoto. Few-Body Syst., 54(7-10):1353–1356, 2013, Study of Tensor Correlations in ^4He via the $^4\text{He}(p, dp)d$ and $^4\text{He}(p, dp)pn$ Reactions.*
- [117] *N Miyawaki, M Fukuda, S Kurashima, H Kashiwagi, S Okumura, K Arakawa, and T Kamiya. Nucl. Instruments Methods Phys. Res. Sect. A Accel. Spectrometers, Detect. Assoc. Equip., 715:126–131, 2013, Geometric analysis of phase bunching in the central region of cyclotron.*
- [118] *A I Morales, J Benlliure, M Górska, H Grawe, S Verma, P H Regan, Z Podolyák, S Pietri, R Kumar, E Casarejos, A Algora, N Alkhomashi, H Álvarez Pol, G Benzoni, A Blazhev, P Boutachkov, A M Bruce, L S Cáceres, I J Cullen, A M Denis Bacelar, P Doornenbal, M E Estévez-Aguado, G Farrelly, Y Fujita, A B Garnsworthy, W Gelletly, J Gerl, J Grebosz, R Hoischen, I Kojouharov, N Kurz, S Lalkowski, Z Liu, C Mihai, F Molina, D Mücher, W Prokopowicz, B Rubio, H Schaffner, S J Steer, A Tamii, S Tashenov, J J Valiente-Dobón, P M Walker, H J Wollersheim, and P J Woods. Phys. Rev. C - Nucl. Phys., 88(1), 2013, β -delayed γ -ray spectroscopy of $^{203,204}\text{Au}$ and $^{200-202}\text{Pt}$.*
- [119] *M Moritsu, S Adachi, M Agnello, S Ajimura, K Aoki, H C Bhang, B Bassalleck, E Botta, S Bufalino, N Chiga, H Ekawa, P Evtoukhovitch, A Feliciello, H Fujioka, S Hayakawa, F Hiruma, R Honda, K Hosomi, Y Ichikawa, M Ieiri, Y Igarashi, K Imai, N Ishibashi, S Ishimoto, K Itahashi, R Iwasaki, C W Joo, S Kanatsuki, M J Kim, S J Kim, R Kiuchi, T Koike, Y Komatsu, V V Kulikov, S Marcello, S Masumoto, Y Matsumoto, K Matsuoka, K Miwa, T Nagae, M Naruki, M Niiyama, H Noumi, Y Nozawa, R Ota, K Ozawa, N Saito, A Sakaguchi, H Sako, V Samoïlov, M Sato, S Sato, Y Sato, S Sawada, M Sekimoto, K Shirotori, H Sugimura, S Suzuki, H Takahashi, T Takahashi, T N Takahashi, H Tamura, T Tanaka, K Tanida, A O Tokiyasu, N Tomida, Z Tsamalaidze, M Ukai, K Yagi, T O Yamamoto, S B Yang, Y Yonemoto, C J Yoon, and K Yoshida. Nucl. Phys. A, 914:91–96, 2013, Search for the $\Theta +$ pentaquark at J-PARC.*

- [120] *N Muramatsu. Few-Body Syst.*, 54(7-10):997–1003, 2013, Recent Progress and Results of LEPS, LEPS2, and ELPH.
- [121] *M Naruki, S Adachi, M Agnello, S Ajimura, K Aoki, H Bang, B Bassalleck, E Botta, S Bufalino, H Ekawa, P Evtoukhovitch, A Feliciello, H Fujioka, S Hayakawa, F Hiruma, R Honda, K Hosomi, Y Ichikawa, M Ieiri, Y Igarashi, K Imai, N Ishibashi, S Ishimoto, K Itahashi, R Iwasaki, G G Joo, S Kanatsuki, M J Kim, S J Kim, R Kiuchi, T Koike, Y Komatsu, V V Kulikov, S Marcello, S Masumoto, Y Matsumoto, K Matsuoka, K Miwa, M Moritsu, T Nagae, M Niiyama, H Noumi, Y Nozawa, R Ota, K Ozawa, N Saito, A Sakaguchi, H Sako, V Samoïlov, M Sato, S Sato, Y Sato, S Sawada, M Sekimoto, K Shirotori, H Sugimura, S Suzuki, T N Takahashi, H Takahashi, T Takahashi, H Tamura, T Tanaka, K Tanida, A O Tokiyasu, N Tomida, Z Tsamalaidze, M Ukai, K Yagi, T O Yamamoto, S B Yang, Y Yonemoto, C J Yoon, and K Yoshida. Few-Body Syst.*, 54(7-10):955–960, 2013, Search for Pentaquark Θ^+ in Hadronic Reaction at J-PARC.
- [122] *M Niiyama. Nucl. Phys. A*, 914:543–552, 2013, Recent results from LEPS and prospects of LEPS II at SPring-8.
- [123] *H Noumi. Few-Body Syst.*, 54(7-10):813–819, 2013, Hadron Experimental Facility at J-PARC.
- [124] *I Ogawa, R Hazama, K Mukaida, K Kishimoto, T Kobayashi, S Tomii, H Sakai, A Katsuki, T Itamura, Y Tanaka, S Umehara, S Yoshida, K Matsuoka, and T Kishimoto. Nucl. Instruments Methods Phys. Res. Sect. A Accel. Spectrometers, Detect. Assoc. Equip.*, 705:1–6, 2013, Background reduction using single-photoelectron counting for WIMP search.
- [125] *H J Ong, I Tanihata, A Tamii, T Myo, K Ogata, M Fukuda, K Hirota, K Ikeda, D Ishikawa, T Kawabata, H Matsubara, K Matsuta, M Mihara, T Naito, D Nishimura, Y Ogawa, H Okamura, A Ozawa, D Y Pang, H Sakaguchi, K Sekiguchi, T Suzuki, M Taniguchi, M Takashina, H Toki, Y Yasuda, M Yosoi, and J Zenihiro. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 725(4-5):277–281, 2013, Probing effect of tensor interactions in ^{16}O via (p,d) reaction.
- [126] *D Patel, U Garg, M Fujiwara, T Adachi, H Akimune, G P A Berg, M N Harakeh, M Itoh, C Iwamoto, A Long, J T Matta, T Murakami, A Okamoto, K Sault, R Talwar, M Uchida, and M Yosoi. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 726(1-3):178–181, 2013, Testing the mutually enhanced magicity effect in nuclear incompressibility via the giant monopole resonance in the $^{204,206,208}\text{Pb}$ isotopes.
- [127] *C Rappold, E Kim, D Nakajima, T R Saito, O Bertini, S Bianchin, V Bozkurt, M Kavatsyuk, Y Ma, F Maas, S Minami, B Özel Tashenov, K Yoshida, P Achenbach, S Ajimura, T Aumann, C Ayerbe Gayoso, H C Bhang, C Caesar, S Erturk, T Fukuda, B Göküzüm, E Guliev, T Hiraiwa, J Hoffmann, G Ickert, Z S Ketenci, D Khanefit, M Kim, S Kim, K Koch, N Kurz, A Le Fèvre, Y Mizoi, M Moritsu, T Nagae, L Nungesser, A Okamura, W Ott, J Pochodzalla, A Sakaguchi, M Sako, C J Schmidt, M Sekimoto, H Simon, H Sugimura, T Takahashi, G J Tambave, H Tamura, W Trautmann, S Voltz, N Yokota, and C J Yoon. Nucl. Phys. A*, 913:170–184, 2013, Hypernuclear spectroscopy of products from ^6Li projectiles on a carbon target at 2A GeV.
- [128] *C Rappold, E Kim, T R Saito, O Bertini, S Bianchin, V Bozkurt, M Kavatsyuk, Y Ma, F Maas, S Minami, D Nakajima, B Özel Tashenov, K Yoshida, P Achenbach, S Ajimura, T Aumann, C Ayerbe Gayoso, H C Bhang, C Caesar, S Erturk, T Fukuda, B Göküzüm, E Guliev, J Hoffmann, G Ickert, Z S Ketenci, D Khanefit, M Kim, S Kim, K Koch,*

N Kurz, A Le Fèvre, Y Mizoi, L Nungesser, W Ott, J Pochodzalla, A Sakaguchi, C J Schmidt, M Sekimoto, H Simon, T Takahashi, G J Tambave, H Tamura, W Trautmann, S Voltz, and C J Yoon. Phys. Rev. C - Nucl. Phys., 88(4), 2013, Search for evidence of $\Lambda^3\text{n}$ by observing $\text{d}+\pi^-$ and $\text{t}+\pi^-$ final states in the reaction of $6\text{Li}+^{12}\text{C}$ at 2A GeV.

- [129] *K Sagara, S Kuroita, T Sueta, H Shimoda, Y Eguchi, K Yashima, T Yabe, M Dozono, Y Yamada, T Wakasa, T Noro, H Matsubara, J Zenihiro, Y Tameshige, H Okamura, A Tamii, K Hatanaka, T Saito, Y Maeda, and H Kamada. Few-Body Syst.*, 54(1-4):469–473, 2013, Cross Section Enhancement in pd Reactions at Higher Energy.
- [130] *S Sakaguchi, Y Iseri, T Uesaka, M Tanifuji, N Aoi, E Hiyama, Y Ichikawa, S Ishikawa, K Itoh, M Itoh, H Iwasaki, T Kawabata, T Kawahara, H Kuboki, Y Maeda, T Nakao, H Okamura, H Sakai, Y Sasamoto, M Sasano, Y Satou, K Sekiguchi, K Suda, D Suzuki, A Tamii, T Wakui, K Yako, M Yamaguchi, and Y Yamamoto. Few-Body Syst.*, 54(7-10):1393–1398, 2013, Analyzing Power in Elastic Scattering of Polarized Protons from Neutron-rich Helium Isotopes.
- [131] *S Sakaguchi, T Uesaka, N Aoi, Y Ichikawa, K Itoh, M Itoh, T Kawabata, T Kawahara, Y Kondo, H Kuboki, T Nakamura, T Nakao, Y Nakayama, H Sakai, Y Sasamoto, K Sekiguchi, T Shimamura, Y Shimizu, and T Wakui. Phys. Rev. C - Nucl. Phys.*, 87(2), 2013, Shallow and diffuse spin-orbit potential for proton elastic scattering from neutron-rich helium isotopes at 71 MeV/nucleon.
- [132] *Y Satou, K Tshoo, H Bhang, S Choi, J W Hwang, T Nakamura, Y Kondo, Y Nakayama, N Kobayashi, K N Tanaka, S Deguchi, Y Kawada, N Tanaka, T Motobayashi, H Sakurai, H Otsu, N Aoi, S Takeuchi, K Yoneda, Y Togano, M Ishihara, S Shimoura, T Kobayashi, M Matsushita, T Honda, T Sumikama, Y Miyashita, K Yoshinaga, N A Orr, F M Marques, J Gibelin, F Delaunay, D Sohler, T Zheng, Z H Li, and Z X Cao. Few-Body Syst.*, 54(1-4):287–290, 2013, Invariant Mass Spectroscopy of ^{23}O via the (p, p^-) Reaction in Inverse Kinematics.
- [133] *Y Shimbara, Y Fujita, T Adachi, G P A Berg, H Fujimura, H Fujita, K Fujita, K Hara, K Y Hara, K Hatanaka, J Kamiya, K Katori, T Kawabata, K Nakanishi, G Martínez-Pinedo, N Sakamoto, Y Sakemi, Y Shimizu, Y Tameshige, M Uchida, M Yoshifuku, and M Yosoi. Phys. Rev. C - Nucl. Phys.*, 88(5), 2013, Erratum: High-resolution study of Gamow-Teller transitions with the $^{37}\text{Cl}(^3\text{He}, \text{t})^{37}\text{Ar}$ reaction (Physical Review C - Nuclear Physics (2012) 86 (024312)).
- [134] *K Shirotori. Few-Body Syst.*, 54(1-4):145–150, 2013, Search for Θ^+ via the Pion Induced Reaction at J-PARC: J-PARC E19.
- [135] *D Steppenbeck, S Takeuchi, N Aoi, P Doornenbal, J Lee, M Matsushita, H Wang, H Baba, N Fukuda, S Go, M Honma, K Matsui, S Michimasa, T Motobayashi, D Nishimura, T Otsuka, H Sakurai, Y Shiga, P.-A. Söderström, T Sumikama, H Suzuki, R Taniuchi, Y Utsuno, J J Valiente-Dobón, and K Yoneda. 445(1)*, 2013, Investigating the strength of the $N = 34$ subshell closure in ^{54}Ca .
- [136] *D Steppenbeck, S Takeuchi, N Aoi, P Doornenbal, M Matsushita, H Wang, H Baba, N Fukuda, S Go, M Honma, J Lee, K Matsui, S Michimasa, T Motobayashi, D Nishimura, T Otsuka, H Sakurai, Y Shiga, P.-A. Söderström, T Sumikama, H Suzuki, R Taniuchi, Y Utsuno, J J Valiente-Dobón, and K Yoneda. Nature*, 502(7470):207–210, 2013, Evidence for a new nuclear 'magic number' from the level structure of ^{54}Ca .
- [137] *H Sugimura, S Ajimura, H C Bhang, L Busso, M Endo, D Faso, T Fukuda, S Hasegawa, S Hayakawa, R Honda, Y Ichikawa, K Imai, N Ishibashi, T Kishimoto, R Kiuchi, K Matsuda, K Matsuoka, K Miwa, Y Mizoi, O Morra, S Nagamiya, M Nakagawa, H Noumi,*

R Ota, P K Saha, H Sako, C Samanta, S Sato, Y Shimizu, K Shirotori, T Soyama, A Sakaguchi, T Takahashi, T N Takahashi, T Tanaka, and K Yoshida. Few-Body Syst., 54(7-10):1235–1238, 2013, Study of Neutron-Rich Hypernuclei by the (π^- , K^+) Reaction at J-PARC.

- [138] *T Sumita, K Morimoto, D Kaji, H Haba, K Ozeki, R Sakai, A Yoneda, A Yoshida, H Hasebe, K Katori, N Sato, Y Wakabayashi, S.-I. Mitsuoka, S.-I. Goto, M Murakami, Y Kariya, F Tokanai, K Mayama, M Takeyama, T Moriya, E Ideguchi, T Yamaguchi, H Kikunaga, J Chiba, and K Morita. J. Phys. Soc. Japan*, 82(2), 2013, New result on the production of ^{277}Cn by the $^{208}\text{Pb} + ^{70}\text{Zn}$ Reaction.
- [139] *H Suzuki, N Aoi, E Takeshita, S Takeuchi, S Ota, H Baba, S Bishop, T Fukui, Y Hashimoto, E Ideguchi, K Ieki, N Imai, M Ishihara, H Iwasaki, S Kanno, Y Kondo, T Kubo, K Kurita, K Kusaka, T Minemura, T Motobayashi, T Nakabayashi, T Nakamura, T Nakao, M Niikura, T Okumura, T K Ohnishi, H J Ong, H Sakurai, S Shimoura, R Sugo, D Suzuki, M K Suzuki, M Tamaki, K Tanaka, Y Togano, and K Yamada. Phys. Rev. C - Nucl. Phys.*, 88(2), 2013, Collectivity of neutron-rich Ti isotopes.
- [140] *H Suzuki, T Kubo, N Fukuda, N Inabe, D Kameda, H Takeda, K Yoshida, K Kusaka, Y Yanagisawa, M Ohtake, H Sato, Y Shimizu, H Baba, M Kurokawa, T Ohnishi, K Tanaka, O B Tarasov, D Bazin, D J Morrissey, B M Sherrill, K Ieki, D Murai, N Iwasa, A Chiba, Y Ohkoda, E Ideguchi, S Go, R Yokoyama, T Fujii, D Nishimura, H Nishibata, S Momota, M Lewitowicz, G DeFrance, I Celikovic, and K Steiger. Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms*, 317(PART B):756–768, 2013, Production cross section measurements of radioactive isotopes by BigRIPS separator at RIKEN RI Beam Factory.
- [141] *A Tamii. Acta Phys. Pol. B*, 44(3):571–580, 2013, Studies of electric dipole response in nuclei using the scattering of polarized protons.
- [142] *A Tamii, N Aoi, H Fujita, Y Fujita, K Hatanaka, T Hashimoto, T Kawabata, K Miki, M Itoh, T Itoh, M Kamimura, K Ogata, H J Ong, H Sakaguchi, T Shima, T Suzuki, and T Yamamoto. Few-Body Syst.*, 54(7-10):1607–1610, 2013, Non-resonant Triple- α Reaction Rate at Low Temperature.
- [143] *H Tamura, K Hosomi, S Bufalino, N Chiga, P Evtoukhovitch, A Feliciello, R Honda, T Koike, Y Ma, K Miwa, A Sasaki, Y Sasaki, K Shirotori, K Tanida, Z Tsamalaidze, M Ukai, T O Yamamoto, and S B Yang. Nucl. Phys. A*, 914:99–108, 2013, Gamma-ray spectroscopy of hypernuclei - present and future.
- [144] *M Tanaka, H Kohri, T Ohta, M Yosoi, M Fujiwara, K Ueda, S Imoto, K Takamatsu, J.-P. Didélez, G Frossati, A de Waard, Y Kiselev, S Makino, H Fujimura, K Fukushima, S Fukushima, and H Kondoh. Phys. Part. Nucl.*, 44(6):959–963, 2013, Production of hyperpolarized nuclei for MRI.
- [145] *I Tanihata. Phys. Scr.*, (T152), 2013, Effects of tensor forces in nuclei.
- [146] *I Tanihata, H Savajols, and R Kanungo. Prog. Part. Nucl. Phys.*, 68(1):215–313, 2013, Recent experimental progress in nuclear halo structure studies.
- [147] *K Tshoo, Y Satou, H Bhang, S Choi, T Nakamura, S Deguchi, Y Kawada, Y Kondo, N Kobayashi, Y Nakayama, K N Tanaka, N Tanaka, N Aoi, M Ishihara, T Motobayashi, H Otsu, H Sakurai, S Takeuchi, Y Togano, K Yoneda, Z H Li, F Delaunay, J Gibelin, F M Marques, N A Orr, T Honda, M Matsushita, T Kobayashi, Y Miyashita, T Sumikama, K Yoshinaga, S Shimoura, D Sohler, T Zheng, and Z X Cao. Few-Body Syst.*, 54(1-4):459–463, 2013, The $N = 16$ Spherical Shell Closure in ^{24}O .

- [148] *H Ueda, M Fukuda, K Hatanaka, T Wang, A Ishiyama, and S Noguchi. IEEE Trans. Appl. Supercond.*, 23(3), 2013, Spatial and temporal behavior of magnetic field distribution due to shielding current in HTS coil for cyclotron application.
- [149] *H Ueda, M Fukuda, K Hatanaka, T Wang, X Wang, A Ishiyama, S Noguchi, S Nagaya, N Kashima, and N Miyahara. IEEE Trans. Appl. Supercond.*, 23(3), 2013, Conceptual design of next generation HTS cyclotron.
- [150] *H Ueno, H Miyatake, Y Yamamoto, S Tanimoto, T Shimoda, N Aoi, K Asahi, E Ideguchi, M Ishihara, H Izumi, T Kishida, T Kubo, S Mitsuoka, Y Mizoi, M Notani, H Ogawa, A Ozawa, M Sasaki, T Shirakura, N Takahashi, and K Yoneda. Phys. Rev. C - Nucl. Phys.*, 87(3), 2013, β -Delayed neutron and γ -ray spectroscopy of ^{17}C utilizing spin-polarized ^{17}B .
- [151] *M Utsuro, T Kuwai, T Ohta, H Kohri, and M Fujiwara. Phys. B Condens. Matter*, 418:36–40, 2013, Anisotropic thermal conductivity of hydrogen deuteride crystal for polarized proton target development.
- [152] *H Wang, N Aoi, S Takeuchi, M Matsushita, P Doornenbal, T Motobayashi, D Steppenbeck, K Yoneda, H Baba, L Cáceres, Zs. Dombrádi, K Kobayashi, Y Kondo, J Lee, K Li, H Liu, R Minakata, D Nishimura, H Otsu, S Sakaguchi, H Sakurai, H Scheit, D Sohler, Y Sun, Z Tian, R Tanaka, Y Togano, Zs. Vajta, Z Yang, T Yamamoto, Y Ye, and R Yokoyama. Phys. Rev. C - Nucl. Phys.*, 88(5), 2013, Collectivity evolution in the neutron-rich Pd isotopes toward the N=82 shell closure.
- [153] *H Wang, N Aoi, S Takeuchi, M Matsushita, P Doornenbal, T Motobayashi, D Steppenbeck, K Yoneda, K Kobayashi, J Lee, H.-N. Liu, Y Kondo, R Yokoyama, H Sakurai, and Y.-L. Ye. Chinese Phys. Lett.*, 30(4), 2013, Observation of new isotope ^{131}Ag via the two-step fragmentation technique.
- [154] *J Yasuda, T Wakasa, M Okamoto, M Dozono, K Hatanaka, M Ichimura, S Kuroita, Y Maeda, T Noro, Y Sakemi, M Sasano, and K Yako. Prog. Theor. Exp. Phys.*, 2013(6), 2013, Extraction of anti-analog giant dipole resonance and neutron skin thickness for ^{208}Pb .

- [1] *M Brenna, X Roca-Maza, G Colò, P F Bortignon, K Mizuyama, and G Pozzi. Phys. Scr., 87(T154), 2013, Low-lying dipole response in stable and unstable nuclei.*
- [2] *A Doté, T Inoue, and T Myo. Nucl. Phys. A, 912:66–101, 2013, Comprehensive application of a coupled-channel complex scaling method to the K-N- π Y system.*
- [3] *T Fukui, K Ogata, K Minomo, M Yahiro, and M Yahiro. Determination of $8\text{B}(p,\gamma)9\text{C}$ reaction rate from 9C breakup. In *Few-Body Syst.*, Vol. 54, pp. 199–204, Research Center for Nuclear Physics, Osaka University, Osaka 567-0047, Japan, 2013. CERN.*
- [4] *H Guo, Y Watanabe, T Matsumoto, K Ogata, and M Yahiro. Phys. Rev. C - Nucl. Phys., 87(2), 2013, Systematic analysis of nucleon scattering from $6,7\text{Li}$ with the continuum discretized coupled channels method.*
- [5] *Sh. Hamada, Y Hirabayashi, N Burtebayev, and S Ohkubo. Phys. Rev. C - Nucl. Phys., 87(2), 2013, Observation of an Airy minimum in elastic and inelastic scattering of 3He from 12C at 50.5 and 60 MeV, and α -particle condensation in 12C .*
- [6] *Y Hirabayashi and S Ohkubo. Phys. Rev. C - Nucl. Phys., 88(1), 2013, Unification of Airy structure in inelastic $\alpha+16\text{O}$ scattering and α -cluster structure with core excitation in 20Ne .*
- [7] *K Horii, H Toki, T Myo, and K Ikeda. Few-Body Syst., 54(1-4):441–445, 2013, Tensor-Optimized Few-Body Model for s-Shell Nuclei.*
- [8] *K Horii, H Toki, T Myo, and K Ikeda. Few-Body Syst., 54(7-10):1367–1370, 2013, Tensor Optimized Few-body Model for Light Nuclei.*
- [9] *A Hosaka, T Hyodo, D Jido, H Nagahiro, K Nawa, S Ohkoda, S Ozaki, Y Yamaguchi, and S Yasui. Few-Body Syst., 54(1-4):19–24, 2013, Composite and Elementary Components in Hadron Resonances.*
- [10] *N Itagaki, T Ichikawa, J A Maruhn, S Ohkubo, T Kokalova, and W V Oertzen. 420(1), 2013, Exotic cluster structure in light nuclei.*
- [11] *H Kamano. Phys. Rev. C - Nucl. Phys., 88(4), 2013, Impact of πn $\pi\pi N$ data on determining high-mass nucleon resonances.*
- [12] *H Kamano, S X Nakamura, T.-S.H. Lee, and T Sato. Phys. Rev. C - Nucl. Phys., 88(3), 2013, Nucleon resonances within a dynamical coupled-channels model of πn and γn reactions.*
- [13] *K Kato, T Myo, Y Kikuchi, and T Yoshida. 436(1), 2013, Multi-cluster problems: Resonances, scattering and condensed states.*
- [14] *K P Khemchandani, A Martínez Torres, H Kaneko, H Nagahiro, and A Hosaka. Few-Body Syst., 54(1-4):343–346, 2013, Resonances Generated by the Vector Meson-Baryon Dynamics.*
- [15] *K P Khemchandani, A Martínez Torres, H Nagahiro, and A Hosaka. Nucl. Phys. A, 914:300–304, 2013, Λ and σ resonances coupled to vector and pseudoscalar mesons.*

- [16] *K P Khemchandani, A Martínez Torres, H Nagahiro, and A Hosaka. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 88(11), 2013, Role of vector and pseudoscalar mesons in understanding 1/2- N^* and Δ resonances.
- [17] *Y Kikuchi, T Matsumoto, K Minomo, and K Ogata. Phys. Rev. C - Nucl. Phys.*, 88(2), 2013, Two neutron decay from the 21+ state of 6He.
- [18] *Y Kikuchi, T Myo, K Kato, and K Ikeda. Few-Body Syst.*, 54(7-10):1445–1448, 2013, Coulomb Breakup Reactions of Two-Neutron Halo Nuclei 6He and 11Li Using Complex-Scaled Solutions of the Lippmann-Schwinger Equation.
- [19] *Y Kikuchi, T Myo, K Kato, and K Ikeda. 436(1)*, 2013, Mechanisms of Coulomb breakup reactions of 6He and 11Li.
- [20] *Y Kikuchi, T Myo, K Kato, and K Ikeda. Phys. Rev. C - Nucl. Phys.*, 87(3), 2013, Coulomb breakup reactions of 11Li in the coupled-channel 9Li+n+n model.
- [21] *S.-H. Kim, S.-I. Nam, A Hosaka, and H.-C. Kim. Few-Body Syst.*, 54(7-10):1499–1502, 2013, Contribution of N^* and Δ^* Resonances in $K^*\Sigma$ (1190) Photoproduction.
- [22] *S.-H. Kim, S.-I. Nam, A Hosaka, and H.-C. Kim. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 88(5), 2013, $K^*\Sigma$ photoproduction off the proton target with baryon resonances.
- [23] *M A Klatt, T Ichikawa, K Iida, N Itagaki, J A Maruhn, K Matsuyanagi, K Mecke, S Ohkubo, P.-G. Reinhard, and B Schuetrumpf. 445(1)*, 2013, Exotic cluster structures in the mean-field theory.
- [24] *N Kurihara, Y Kikuchi, and K Kato. Few-Body Syst.*, 54(7-10):1381–1384, 2013, Study of the Isospin Mixing in 6Li with $\alpha + p + n$ Three-Body Model.
- [25] *A Martínez Torres, K P Khemchandani, D Jido, Y Kanada-En'yo, and A Hosaka. Few-Body Syst.*, 54(1-4):333–337, 2013, Few Body Systems Made of Pseudoscalars.
- [26] *T Matsumoto, K Minomo, K Ogata, M Yahiro, and K Kato. Few-Body Syst.*, 54(7-10):1437–1440, 2013, Four-Body CDCC Analysis for Breakup Reactions of Three-Body Projectiles.
- [27] *K Minomo, T Matsumoto, K Ogata, and M Yahiro. Few-Body Syst.*, 54(7-10):1417–1419, 2013, Eikonal Reaction Theory for One- and Two-Neutron Removal Reactions.
- [28] *R Molina, H Nagahiro, A Hosaka, T Branz, and E Oset. Few-Body Syst.*, 54(7-10):1259–1262, 2013, Dynamically Generated Resonances from Two Vectors in the Charm Sector and their Decays.
- [29] *T Myo. Few-Body Syst.*, 54(7-10):849–855, 2013, Role of Tensor Force in Light Nuclei with Tensor-Optimized Shell Model.
- [30] *T Myo, Y Kikuchi, and K Kato. Phys. Rev. C - Nucl. Phys.*, 87(4), 2013, Erratum: Five-body resonances of 8C using the complex scaling method (Physical Review C - Nuclear Physics (2012) 85 (034338)).
- [31] *T Myo, A Umeya, H Toki, and K Ikeda. 436(1)*, 2013, Role of the tensor force in light nuclei with the tensor-optimized shell model.
- [32] *T Myo, A Umeya, H Toki, and K Ikeda. Few-Body Syst.*, 54(1-4):451–454, 2013, Role of Tensor Force in He and Li Isotopes with Tensor Optimized Shell Model.
- [33] *H Nagahiro and A Hosaka. Phys. Rev. C - Nucl. Phys.*, 88(5), 2013, Composite and elementary nature of a resonance in the σ model.

- [34] *K Nawa, S Ozaki, H Nagahiro, D Jido, and A Hosaka. Prog. Theor. Exp. Phys.*, 2013(8), 2013, Complex 2D matrix model and geometrical map on the complex-Nc plane.
- [35] *K Ogata, T Myo, T Furumoto, T Matsumoto, and M Yahiro. Phys. Rev. C - Nucl. Phys.*, 88(2), 2013, Interplay between the 02+ resonance and the nonresonant continuum of the drip-line two-neutron halo nucleus 22C.
- [36] *S Ohkoda, Y Yamaguchi, S Yasui, K Sudoh, and A Hosaka. Few-Body Syst.*, 54(7-10):1019–1022, 2013, Exotic Mesons with Hidden Bottom Near Thresholds.
- [37] *S Ohnishi, Y Ikeda, H Kamano, and T Sato. Phys. Rev. C - Nucl. Phys.*, 88(2), 2013, Signature of strange dibaryons in kaon- and photon-induced reactions.
- [38] *S Ohnishi, Y Ikeda, H Kamano, and T Sato. Few-Body Syst.*, 54(7-10):1119–1122, 2013, Production Reaction of KNN- π YN Resonance from Faddeev Equations.
- [39] *S Ohnishi, Y Ikeda, H Kamano, and T Sato. Few-Body Syst.*, 54(1-4):347–351, 2013, Signature of Strange Dibaryon in Kaon-Induced Reaction.
- [40] *T Sakuda and S Ohkubo. Nucl. Phys. A*, 908:73–76, 2013, Cluster structure and deformed bands in the 38Ar nucleus.
- [41] *S Sasabe, T Matsumoto, S Tagami, N Furutachi, K Minomo, Y R Shimizu, and M Yahiro. Phys. Rev. C - Nucl. Phys.*, 88(3), 2013, Reaction mechanism in odd-even staggering of reaction cross sections.
- [42] *A Umeya, T Myo, E Hiyama, H Toki, and K Ikeda. Few-Body Syst.*, 54(1-4):99–102, 2013, s-Shell Λ -Hypernuclei Taking into Account Λ N- Σ N Coupling Explicitly with TOSM.
- [43] *A Umeya, T Myo, E Hiyama, H Toki, and K Ikeda. Few-Body Syst.*, 54(7-10):1215–1217, 2013, Tensor-Optimized Shell Model for s-Shell Hypernuclei.
- [44] *Y Yamaguchi, S Ohkoda, S Yasui, and A Hosaka. Phys. Rev. D - Part. Fields, Gravit. Cosmol.*, 87(7), 2013, Hadronic molecules for charmed and bottom baryons near thresholds.
- [45] *Y Yamaguchi, S Ohkoda, S Yasui, and A Hosaka. Few-Body Syst.*, 54(7-10):1051–1054, 2013, Exotic Baryons from a Heavy Meson and a Nucleon.
- [46] *S Yasui, S Ohkoda, Y Yamaguchi, K Sudoh, and A Hosaka. Few-Body Syst.*, 54(7-10):1023–1026, 2013, Doubly Charmed Exotic Mesons.
- [47] *S Yasui, K Sudoh, Y Yamaguchi, S Ohkoda, A Hosaka, and T Hyodo. Phys. Lett. Sect. B Nucl. Elem. Part. High-Energy Phys.*, 727(1-3):185–189, 2013, Spin degeneracy in multi-hadron systems with a heavy quark.

- [1] T Yorita, M Fukuda, K Hatanaka, K Kamakura, Y Yasuda, S Morinobu, and A Tamii. Study of extraction and transport of intense highly charged ions for 18GHZ SC-ECRIS at RCNP. pp. 145–147, RCNP, Osaka, Japan, 2013.
- [2] T Yorita, N Muramatsu, T Hotta, T Nakano, M Oka, M Yosoi, S Daté, Y Ohashi, H Ohkuma, M Oishi, Y Okayasu, M Shoji, S Suzuki, and Y Taniuchi. Production of intense high energy gamma beam for LEPS2 project at SPring-8. pp. 2162–2164, RCNP, Osaka, Japan, 2013.
- [3] E Aguayo, F T Avignone III, H O Back, A S Barabash, J R Beene, M Bergevin, F E Bertrand, M Boswell, V Brudanin, M Busch, Y-D. Chan, C D Christofferson, J I Collar, D C Combs, R J Cooper, J A Detwiler, P J Doe, Y Efremenko, V Egorov, H Ejiri, S R Elliott, J Esterline, J E Fast, N Fields, P Finnerty, F M Fraenkle, V M Gehman, G K Giovanetti, M P Green, V E Guiseppe, K Gusey, A L Hallin, R Hazama, R Henning, A Hime, E W Hoppe, M Horton, S Howard, M A Howe, R A Johnson, K J Keeter, C Keller, M F Kidd, A Knecht, O Kochetov, S I Konovalov, R T Kouzes, B D LaFerriere, B H LaRoque, J Leon, L E Leviner, J C Loach, S MacMullin, M G Marino, R D Martin, D.-M. Mei, J H Merriman, M L Miller, L Mizouni, M Nomachi, J L Orrell, N R Overman, D G Phillips II, A W P Poon, G Perumpilly, G Prior, D C Radford, K Rielage, R G H Robertson, M C Ronquest, A G Schubert, T Shima, M Shirchenko, K J Snively, D Steele, J Strain, K Thomas, V Timkin, W Tornow, I Vanyushin, R L Varner, K Vetter, K Vorren, J F Wilkerson, E Yakushev, A R Young, C.-H. Yu, V I Yumatov, and C Zhang. The Majorana double beta decay experiment: Present status. pp. 164–168, Department of Physics, Black Hills State University, Spearfish, SD, United States, 2013.
- [4] K Hatanaka. RCNP cyclotron facility and application program. Vol. 1533, pp. 17–24, Research Center for Nuclear Physics, Osaka University, Mihogaoka 10-1, Ibaraki, Osaka, 567-0047, Japan, 2013.
- [5] K Hatanaka, M Fukuda, N Hamatani, K Kamakura, T Saito, H Ueda, T Yorita, Y Yasuda, and T Kawaguchi. HTS magnet to polarize ultra cold neutrons. pp. 3522–3524, Research Center for Nuclear Physics, Osaka University, 10-1 Mihogaoka, Ibaraki, Osaka, 567-0047, Japan, 2013.
- [6] T Inoue, H Arikawa, S Ezure, K Harada, T Hayamizu, T Ishikawa, M Itoh, T Kato, H Kawamura, H S Nataraj, T Sato, T Aoki, T Furukawa, A Hatakeyama, K Hatanaka, K Imai, T Murakami, Y Shimizu, T Wakasa, H P Yoshida, and Y Sakemi. Search for electron EDM with laser cooled radioactive atom. Vol. 1533, pp. 134–139, Cyclotron and Radioisotope Center, Tohoku University, Miyagi 980-8578, Japan, 2013.
- [7] K Matsuta, Y Masuda, K Hatanaka, S Kawasaki, R Matsumiya, M Mihara, S C Jeong, Y Watanabe, T Adachi, D Nishimura, K Asahi, J W Martin, A Konaka, A Miller, C Bidinosti, T Dawson, L Lee, C A Davis, W D Ramsay, W T H Van Oers, E Korkmaz, and L Buckman. An EDM measurement with a new comagnetometer and a high density UCN source. Vol. 1560, pp. 152–154, Dept. of Physics, Osaka Univ., Toyonaka Osaka560-0043, Japan, 2013.
- [8] S Umehara, T Kishimoto, M Nomachi, S Yoshida, S Ajimura, K Suzuki, K Ichimura, K Matsuoka, N Nakatani, G Ito, H Kakubata, M Saka, W Wang, J Takemoto, W Chan,

M Doihara, Y Tamagawa, I Ogawa, S Ueno, S Maeda, A Yamamoto, S Tomita, G Fujita, A Kawamura, T Harada, K Fushimi, R Hazama, H Ohsumi, and K Okada. Search for neutrino-less double beta decay by CANDLES. Vol. 1533, pp. 115–120, Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 560-0047, Japan, 2013.

- [9] H Utsunomiya, T Shima, K Takahisa, D M Filipescu, O Tesileanu, I Gheorghe, H.-T. Nyhus, T Renstrom, Y.-W. Lui, Y Kitagawa, S Amano, and S Miyamoto. Energy calibration of the NewSUBARU storage ring by laser compton-scattering gamma rays and its applications. Department of Physics, Konan University, Okamoto 8-9-1, Higashinada, Kobe 658-8501, Japan, 2013. Institute of Electrical and Electronics Engineers Inc.
- [10] T Wakasa, M Okamoto, M Dozono, K Hatanaka, M Ichimura, S Kuroita, Y Maeda, H Miyasako, T Noro, T Saito, Y Sakemi, T Yabe, and K Yako. Decomposing spin-dipole resonances by complete polarization transfer measurements. Vol. 1524, pp. 46–51, Department of Physics, Kyushu University, Higashi, Fukuoka 812-8581, Japan, 2013.

- [1] T Matsumoto, K Minomo, K Ogata, M Yahiro, and K Kato. CDCC analysis for breakup of three-body projectiles. pp. 205–210, Department of Physics, Kyushu University, Fukuoka 812-8581, Japan, 2013. CERN.
- [2] K Ogata, T Myo, T Furumoto, T Matsumoto, and M Yahiro. Breakup of ^{22}C studied by CDCC with Cluster-orbital shell-model wave functions. pp. 211–216, Research Center for Nuclear Physics, Osaka University, Ibaraki 567-0047, Japan, 2013. CERN.
- [3] S Watanabe, T Matsumoto, K Minomo, K Ogata, and M Yahiro. Comparison of breakup processes of ^6He and ^6Li with four-body CDCC. pp. 217–222, Kyushu University, Fukuoka, Japan, 2013. CERN.
- [4] M Yahiro, K Ogata, T Matsumoto, and K Minomo. Microscopic approach to the scattering of unstable nuclei at intermediate incident energies. pp. 45–52, Department of Physics, Kyushu University, Fukuoka 812-8581, Japan, 2013. CERN.
- [5] T Fukui, K Ogata, K Minomo, M Yahiro, and M Yahiroro. Determination of $^8\text{B}(p,\gamma)^9\text{C}$ reaction rate from ^9C breakup. In *Few-Body Syst.*, Vol. 54, pp. 199–204, Research Center for Nuclear Physics, Osaka University, Osaka 567-0047, Japan, 2013. CERN.
- [6] K Minomo, T Matsumoto, K Ogata, and M Yahiro. Eikonal reaction theory for neutron removal reactions. pp. 61–66, Department of Physics, Kyushu University, Fukuoka 812-8581, Japan, Research Center for Nuclear Physics (RCNP), Osaka University, Ibaraki 567-0047, Japan, 2013. CERN.