


Entanglement suppression for $\Omega\Omega$ scatterings



**Tao-Ran Hu, Katsuyoshi Sone,
Feng-Kun Guo, Tetsuo Hyodo, Ian Low**

RCNP, The University of Osaka

2026, Aug. 18th ₁

Basic concept of entanglement suppression

Entanglement power (EP) of S-matrix $E(\hat{S})$

- ability of generating entanglement



$$|\psi_{\text{out}}\rangle = \hat{S} |\psi_{\text{in}}\rangle$$


 $E(\hat{S})$
entangle?



Entanglement suppression

S.R. Beane, D.B. Kaplan, N. Klco, M.J. Savage, PRL122, 102001 (2019)

$E(\hat{S}) = 0 \rightarrow$ emergent **symmetries** of NN scattering

Generalization to other hadrons and nuclei, ...

Q. Liu, I. Low, T. Mehen, PRC107, 025204 (2023);

T. Kirchner, W. Elkamhawy, H.-W. Hammer, Few-Body Syst. 65, 29 (2024);

T.R. Hu, S. Chen, F.K. Guo, PRD110, 014001 (2024); ...

Talks by Skibiński (Mon, Plenary), Kirchner (Mon, III-1), Griebhammer (Mon, III-1)

Entanglement measures

Entanglement **measure** for bipartite **state** $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$

- von Neumann entropy

$$\mathcal{E}_{\text{vN}}(|\psi\rangle) = -\text{Tr}_A[\rho_A \log \rho_A], \quad \rho_A = \text{Tr}_B |\psi\rangle\langle\psi|$$

- linear entropy (expansion $\log \rho_A \sim (\rho_A - 1)$)

$$\mathcal{E}(|\psi\rangle) = 1 - \text{Tr}_A[\rho_A^2]$$

- product state: minimum (=0)



$$|\uparrow\downarrow\rangle = |\uparrow_A\rangle \otimes |\downarrow_B\rangle$$

- Bell state: maximum (=1/2)

$$|\Psi^\pm\rangle = \frac{1}{\sqrt{2}} |\uparrow\downarrow\rangle \pm \frac{1}{\sqrt{2}} |\downarrow\uparrow\rangle$$

Linear combination:

$$|\psi\rangle = \cos \theta |\uparrow\downarrow\rangle + \sin \theta |\downarrow\uparrow\rangle$$



Entanglement power

Entanglement **power** (EP) of **operator** $\hat{U}$ acting on $|\psi\rangle$

P. Zanardi, C. Zalka, L. Faoro, PRA62, 030301 (2000)

- average over entanglement measure of $\hat{U}|\psi_A\rangle \otimes |\psi_B\rangle$

$$E(\hat{U}) = \overline{\mathcal{E}(\hat{U}|\psi_A\rangle \otimes |\psi_B\rangle)}$$

$$= \int d\omega_A d\omega_B \mathcal{E}(\hat{U}|\psi_A(\omega_A)\rangle \otimes |\psi_B(\omega_B)\rangle)$$

$$\mathcal{E}(|\psi_A\rangle \otimes |\psi_B\rangle) = 0$$

—> ability of $\hat{U}$ to **generate** entanglement (state independent)

- **spin 1/2 (qubit)**: rays in $\mathbb{C}^2 \simeq \mathbb{CP}^1$ —> Fubini-Study measure

$$|\psi(\omega_2)\rangle = \begin{pmatrix} \cos \theta_1 \\ \sin \theta_1 e^{i\nu_1} \end{pmatrix}, \quad d\omega_2 = \frac{1}{\pi} d\theta_1 \cos \theta_1 \sin \theta_1 d\nu_1$$

$$\theta_1 \in [0, \pi/2) \\ \nu_1 \in [0, 2\pi)$$

(equivalent to Bloch sphere with $\theta_1 = \theta/2$, $\nu_1 = \phi$, $d\omega_2 = d\Omega/4\pi$)

S-matrix for NN scattering

S-matrix for pn scattering: 2 phase shifts δ_0, δ_1

$$\hat{S} = \underbrace{\exp\{2i\delta_0\}}_{J=0} \frac{1 - \sigma_A \cdot \sigma_B}{4} + \underbrace{\exp\{2i\delta_1\}}_{J=1} \frac{3 + \sigma_A \cdot \sigma_B}{4}$$

Entanglement power of NN scattering

S.R. Beane, D.B. Kaplan, N. Klco, M.J. Savage, PRL122, 102001 (2019)


$$E(\hat{S}) = \frac{1}{6} \sin^2[2(\delta_0 - \delta_1)] \geq 0$$

$E(\hat{S}) = 0$ is achieved if

$$\begin{cases} \delta_0 = \delta_1 & \text{spin-flavor SU(4)} \\ (\delta_0, \delta_1) = (0,0), (0,\pi/2), (\pi/2,0), (\pi/2,\pi/2) \end{cases}$$

$|a_0| \rightarrow \infty$: NR conformal symmetry

$J=1$



$\delta_0 = \delta_1$: $\hat{S} \sim$ Identity gate, $|\delta_0 - \delta_1| = \pi/2$: $\hat{S} \sim$ SWAP gate

I. Low, T. Mehen, PRD104, 074014 (2021)

Further evidences of entanglement suppression

Spin-flavor symmetries with three flavors

S.R. Beane, D.B. Kaplan, N. Klco, M.J. Savage, PRL122, 102001 (2019)

- $SU(6) \sim (u, d, s) \otimes (\uparrow, \downarrow) \leftarrow$ large N_c
- $SU(16) \sim (n, p, \Lambda, \dots) \otimes (\uparrow, \downarrow) \leftarrow$ Entanglement suppression
- Lattice QCD evaluation of LEC favors $SU(16)$

M.L. Wagman, *et al.*, (NPLQCD), PRD96, 114510 (2017)


Momentum dependence of Entanglement power

Q. Liu, I. Low, PLB, 856, 138899 (2024)

- $np, \Lambda p$ scatterings

$$E(p_{\text{CM}}) = \frac{1}{6} \sin^2[2(\delta_0(p_{\text{CM}}) - \delta_1(p_{\text{CM}}))]$$

- EP suppressed at low energies



Motivation

Decuplet baryons

- spin 3/2
- ground states with Octet in quark model

Large scattering length a_0 ($\delta \sim \pi/2$)?


- $\Delta\Delta$ dibaryon, $d^*(2380)$?

F. Dyson, N.H. Xuong, PRL13, 815 (1964);

P. Adlarson *et al.*, (WASA-at-COSY), PRL102, 052301 (2009)

- lattice QCD for $\Omega\Omega$ scattering: $a_0(J=0) = 4.6 \text{ fm} \gg 1 \text{ fm}$

S. Gongyo, *et al.*, (HAL QCD), PRL120, 212001 (2018)



Quantum information viewpoint

- spin 3/2 scattering = 4d-qudit quantum gates

Spin 3/2 state

Spin 3/2 (4d-qudit): rays in $\mathbb{C}^4 \simeq \mathbb{CP}^3$

$$|\psi_4\rangle = \begin{pmatrix} \cos \theta_1 \sin \theta_2 \sin \theta_3 \\ \sin \theta_1 \sin \theta_2 \sin \theta_3 e^{i\nu_1} \\ \cos \theta_2 \sin \theta_3 e^{i\nu_2} \\ \cos \theta_3 e^{i\nu_3} \end{pmatrix}, \quad d\omega_4 = \frac{3!}{\pi^3} \prod_{i=1}^3 d\theta_i d\nu_i \cos \theta_i \sin^{2i-1} \theta_i,$$

$$\theta_i \in [0, \pi/2), \quad \nu_i \in [0, 2\pi)$$

Spin decomposition (2 symmetric, 2 anti-symmetric)

$$\frac{3}{2} \otimes \frac{3}{2} = \underbrace{0 \oplus 2}_A \oplus \underbrace{1 \oplus 3}_S$$

S-matrix (distinguishable): 4 components with $\delta_0, \delta_1, \delta_2, \delta_3$

$$\hat{S} = \mathcal{J}_0 \exp\{2i\delta_0\} + \mathcal{J}_1 \exp\{2i\delta_1\} + \mathcal{J}_2 \exp\{2i\delta_2\} + \mathcal{J}_3 \exp\{2i\delta_3\}$$

$$\mathcal{J}_0 = \frac{33}{128} + \frac{31}{96} (\mathbf{t}_{3/2} \cdot \mathbf{t}_{3/2}) - \frac{5}{72} (\mathbf{t}_{3/2} \cdot \mathbf{t}_{3/2})^2 - \frac{1}{18} (\mathbf{t}_{3/2} \cdot \mathbf{t}_{3/2})^3, \dots$$

Entanglement power

Entanglement power (~3 days on mathematica)

$$\begin{aligned}
 E(\hat{S}) = \frac{1}{200000} \{ & 77482 - 2100 \cos \left[2 (\delta_0 + \delta_1 - \delta_2 - \delta_3) \right] \\
 & - 2100 \cos \left[2 (\delta_0 - \delta_1 + \delta_2 - \delta_3) \right] - 2100 \cos \left[2 (\delta_0 - \delta_1 - \delta_2 + \delta_3) \right] \\
 & - 1200 \cos \left[2 (\delta_0 - 2\delta_1 + \delta_2) \right] - 4200 \cos \left[2 (\delta_0 + \delta_2 - 2\delta_3) \right] \\
 & - 8400 \cos \left[2 (\delta_1 - 2\delta_2 + \delta_3) \right] \\
 & - 375 \cos \left[4 (\delta_0 - \delta_1) \right] - 10800 \cos \left[2 (\delta_0 - \delta_2) \right] - 625 \cos \left[4 (\delta_0 - \delta_2) \right] \\
 & - 875 \cos \left[4 (\delta_0 - \delta_3) \right] - 2175 \cos \left[4 (\delta_1 - \delta_2) \right] - 26376 \cos \left[2 (\delta_1 - \delta_3) \right] \\
 & - 5481 \cos \left[4 (\delta_1 - \delta_3) \right] - 10675 \cos \left[2 (\delta_2 - \delta_3) \right] \}
 \end{aligned}$$

$E(\hat{S}) = 0$ is achieved if (all cos = +1)

$$\delta_0 = \delta_2 = \delta_{\text{even}}, \quad \delta_1 = \delta_3 = \delta_{\text{odd}}, \quad \begin{cases} |\delta_{\text{even}} - \delta_{\text{odd}}| = 0 & : \text{Identity gate} \\ |\delta_{\text{even}} - \delta_{\text{odd}}| = \pi/2 & : \text{SWAP gate} \end{cases}$$

Identical particles and sub-unitary S-matrix

Identical particles: symmetric spin states are forbidden

- nn scattering/ $\Omega\Omega$ scattering: only antisymmetric component

$$\hat{S}_{nn} = \mathcal{I}_0 \exp\{2i\delta_0\} \quad \text{spin 1 absent}$$

$$\hat{S}_{\Omega\Omega} = \mathcal{I}_0 \exp\{2i\delta_{0,28}\} + \mathcal{I}_2 \exp\{2i\delta_{2,28}\} \quad \text{spin 1 and 3 absent}$$

S-matrix is not unitary in full space of $|\psi_{\text{in}}\rangle : \hat{S}_{nn}\hat{S}_{nn}^\dagger = \mathcal{I}_0 \neq 1$

$$\hat{S} = \begin{pmatrix} (\text{Antisymmetric}) & 0 \\ 0 & (\text{Symmetric}) \end{pmatrix}, \quad \mathcal{I}_0 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

EP is obtained by averaging over initial **product states**

$$E(\hat{U}) = \overline{\mathcal{E} \left(\hat{U} |\psi_A\rangle \otimes |\psi_B\rangle \right)}$$

neither antisymmetric nor symmetric

Normalized entanglement power

Naive calculation for nn scattering: larger than **1/2**?

$$E(\hat{S}_{nn}) = \frac{23}{24}$$

Density matrix of out state is not properly normalized

$$\rho = |\psi_{\text{out}}\rangle\langle\psi_{\text{out}}| = \hat{S}_{nn} |\psi_{\text{in}}\rangle\langle\psi_{\text{in}}| \hat{S}_{nn}^\dagger, \quad \text{Tr}[\rho] = \langle\psi_{\text{in}}| \mathcal{I}_0 |\psi_{\text{in}}\rangle < 1$$

—> Properly **normalized** Entanglement power

Normalized density matrix

$$\tilde{\rho} = \frac{|\psi_{\text{out}}\rangle\langle\psi_{\text{out}}|}{\langle\psi_{\text{out}}|\psi_{\text{out}}\rangle} \quad \Rightarrow \quad E(\hat{S}_{nn}, \tilde{\rho}) = \frac{1}{2}$$

- consistent with J=0 final state (Bell state)

$\Omega\Omega$ scattering

$\Omega\Omega$ S-matrix

$$\hat{S}_{\Omega\Omega} = \mathcal{I}_0 \exp\{2i\delta_{0,28}\} + \mathcal{I}_2 \exp\{2i\delta_{2,28}\}$$

Properly normalized EP of $\Omega\Omega$ S-matrix

$$E_2(\hat{S}_{\Omega\Omega}) = \frac{1}{48} \left\{ 25 - \cos \left[4 (\delta_{0,28} - \delta_{2,28}) \right] \right\}$$

$E_2(\hat{S}_{\Omega\Omega})$ is minimized (1/2, not 0) if

$$\begin{cases} |\delta_{0,28} - \delta_{2,28}| = 0 & \hat{S} \propto \mathcal{I}_0 + \mathcal{I}_2 \equiv \mathcal{P}_A & \text{spin SU(4)} \\ |\delta_{0,28} - \delta_{2,28}| = \pi/2 & \hat{S} \propto -\mathcal{I}_0 + \mathcal{I}_2 \equiv \text{SWAP}_A & \text{NR conformal} \end{cases}$$

$a_0(J=0) \gg 1$ fm by lattice QCD suggests either

- $|a_0(J=2)| \gg 1$ fm (**unitary limit**, $|\delta_{0,28} - \delta_{2,28}| = 0$) or
- $a_0(J=2) \sim 0$ (**noninteracting**, $|\delta_{0,28} - \delta_{2,28}| = \pi/2$)

Summary

- Entanglement suppression for spin 3/2
- Final state should be properly **normalized** for identical particle (e.g. $\Omega\Omega$) scattering
- $\Omega\Omega$ scattering **emergent symmetries**
 - spin SU(4) symmetry ($\hat{S} \propto \mathcal{P}_A$)
 - NR conformal ($\hat{S} \propto \text{SWAP}_A$)
- Application**
 - Spin 2 scattering is constrained from spin 0

T.R. Hu, K. Sone, F.K. Guo, T. Hyodo, I. Low, Phys. Rev. Res. 7, 043306 (2025);
K. Sone, T.R. Hu, F.K. Guo, T. Hyodo, I. Low, arXiv:2602.09630 [hep-ph]